

Title:

HYDANTOIN DERIVATIVE COMPOUNDS, PHARMACEUTICAL
COMPOSITIONS, AND METHODS OF USING SAME

Inventor(s): Gregory S. HAMILTON

NATH & ASSOCIATES
1030 Fifteenth Street, N.W.
Sixth Floor
Washington, D.C. 20005-1503
(202) 775-8383

HYDANTOIN DERIVATIVE COMPOUNDS, PHARMACEUTICAL
COMPOSITIONS, AND METHODS OF USING SAME

BACKGROUND OF THE INVENTION

5

Field of Invention

This invention relates to novel hydantoin and
hydantoin derivative compounds, their inclusion in
pharmaceutical compositions, and their preparation and use
10 for preventing and/or treating neurological disorders; for
treating alopecia and promoting hair growth; for treating
vision disorders and/or improving vision; for treating
memory impairment and/or enhancing memory performance; and
for preventing and/or treating hearing loss in an animal.

15

Description of Related Art

It has been found that picomolar concentrations of an
immunosuppressant such as FK506 and rapamycin stimulate
neurite outgrowth in PC12 cells and sensory nervous, namely
dorsal root ganglion cells (DRGs). Lyons et al., *Proc. Of*
20 *Natl. Acad. Sci.*, 1994 vol. 91, pp. 3191-3195. In whole
animal experiments, FK506 has been shown to stimulate nerve
regeneration following facial nerve injury and results in
functional recovery in animals with sciatic nerve lesions.

Several neurotrophic factors effecting specific
25 neuronal populations in the central nervous system have
been identified. For example, it has been hypothesized
that Alzheimer's disease results from a decrease or loss
of nerve growth factor (NGF). It has thus been proposed
to treat Alzheimer's patients with exogenous nerve growth
30 factor or other neurotrophic proteins such as brain derived
nerve factor (BDNF), glial derived nerve factor, ciliary
neurotrophic factor, and neurotrophin-3 to increase the
survival of degenerating neuronal populations.

Clinical application of these proteins in various
35 neurological disease states is hampered by difficulties in
the delivery and bioavailability of large proteins to

nervous system targets. By contrast, immunosuppressant drugs with neurotrophic activity are relatively small and display excellent bioavailability and specificity. However, when administered chronically, immunosuppressants exhibit a number of potentially serious side effects including nephrotoxicity, such as impairment of glomerular filtration and irreversible interstitial fibrosis (Kopp et al., 1991, *J. Am. Soc. Nephrol.* 1:162); neurological deficits, such as involuntary tremors, or non-specific cerebral angina such as non-localized headaches (De Groen et al., 1987, *N. Engl. J. Med.* 317:861); and vascular hypertension with complications resulting therefrom (Kahan et al., 1989 *N. Engl. J. Med.* 321: 1725).

Accordingly, there is a need for non-immunosuppressive, small-molecule compounds which are useful for neurotrophic effects and for treating neurodegenerative disorders.

Hair loss occurs in a variety of situations. These situations include male pattern alopecia, alopecia senilis, alopecia areata, diseases accompanied by basic skin lesions or tumors, and systematic disorders such as nutritional disorders and internal secretion disorders. The mechanisms causing hair loss are very complicated, but in some instances can be attributed to aging, genetic disposition, the activation of male hormones, the loss of blood supply to hair follicles, and scalp abnormalities.

The immunosuppressant drugs FK506, rapamycin and cyclosporin are well known as potent T-cell specific immunosuppressants, and are effective against graft rejection after organ transplantation. It has been reported that topical, but not oral, application of FK506 (Yamamoto et al., *J. Invest. Dermatol.*, 1994, 102, 160-164; Jiang et al., *J. Invest. Dermatol.* 1995, 104, 523-525) and cyclosporin (Iwabuchi et al., *J. Dermatol. Sci.* 1995, 9,

64-69) stimulates hair growth in a dose-dependent manner. One form of hair loss, alopecia areata, is known to be associated with autoimmune activities; hence, topically administered immunomodulatory compounds are expected to demonstrate efficacy for treating that type of hair loss. The hair growth stimulating effects of FK506 have been the subject of an international patent filing covering FK506 and structures related thereto for hair growth stimulation (Honbo et al., EP 0 423 714 A2). Honbo et al. discloses the use of relatively large tricyclic compounds, known for their immunosuppressive effects, as hair revitalizing agents.

The hair growth and revitalization effects of FK506 and related agents are disclosed in many U.S. patents (Goulet et al., U.S. Patent No. 5,258,389; Luly et al., U.S. Patent No. 5,457,111; Goulet et al., U.S. Patent No. 5,532,248; Goulet et al., U.S. Patent No. 5,189,042; and Ok et al., U.S. Patent No. 5,208,241; Rupprecht et al., U.S. Patent No. 5,284,840; Organ et al., U.S. Patent No. 5,284,877). These patents claim FK506 related compounds. Although they do not claim methods of hair revitalization, they disclose the known use of FK506 for effecting hair growth. Similar to FK506 (and the claimed variations in the Honbo et al. patent), the compounds claimed in these patents are relatively large. Further, the cited patents relate to immunomodulatory compounds for use in autoimmune related diseases, for which FK506's efficacy is well known.

Other U.S. patents disclose the use of cyclosporin and related compounds for hair revitalization (Hauer et al., U.S. Patent No. 5,342,625; Eberle, U.S. Patent No. 5,284,826; Hewitt et al., U.S. Patent No. 4,996,193). These patents also relate to compounds useful for treating autoimmune diseases and cite the known use of cyclosporin and related immunosuppressive compounds for hair growth.

However, immunosuppressive compounds by definition

suppress the immune system and also exhibit other toxic side effects. Accordingly, there is a need for non-immunosuppressive, small molecule compounds which are useful as hair revitalizing compounds.

5 The visual system is composed of the eyes, ocular adnexa and the visual pathways. Dysfunction of the visual system may lead to permanent or temporary visual impairment, i.e. a deviation from normal in one or more
10 in various ways and includes a broad range of visual dysfunctions and disturbances. Without limitation, these dysfunctions and disturbances include partial or total loss of vision, the need for correction of visual acuity for objects near and far, loss of visual field, impaired ocular
15 motility without diplopia (double vision), impaired or skewed color perception, limited adaptation to light and dark, diminished accommodation, metamorphopsic distortion, impaired binocular vision, paresis of accommodation, iridoplegia, entropion, ectropion, epiphora, lagophthalmos,
20 and scarring. See *Physicians' Desk Reference (PDR) for Ophthalmology*, 16th Edition, 6:47 (1988). The visual system may be adversely affected by various ophthalmologic disorders, diseases, injuries, and complications, including, without limitation, genetic disorders; disorders
25 associated with aging or degenerative diseases; disorders correlating to physical injury to the eye, head, or other parts of the body resulting from external forces; disorders resulting from environmental factors; disorders resulting from a broad range of diseases; and combinations of any of
30 the above.

 The visual system is a complex system composed of numerous components. Visual impairment can involve the entire visual system, any one component, or any combination of components, depending upon the precise nature of the

circumstances. The eye is composed of a lens, which is suspended in the zonules of Zinn and is focused by the ciliary body. The ciliary body also secretes aqueous humor, which fills the posterior chamber, passes through
5 the pupil into the anterior chamber, then drains primarily via the canal of Schlemm. The iris regulates the quantity of light entering the eye by adjusting the size of its central opening, the pupil. A visual image is focused onto the retina, the fovea centralis being the retinal area of
10 sharpest visual acuity. The conjunctiva is the mucus membrane which lines the eyelids and the eyeball, and ends abruptly at the limbus conjunctivae, the edge of the conjunctiva overlapping the cornea. The cornea is the clear, transparent anterior portion of the fibrous coat of
15 the eye; it is important in light refraction and is covered with an epithelium that differs in many respects from the conjunctival epithelium.

The retina is the innermost, light sensitive portion of the eye, containing two types of photoreceptors, cones,
20 which are responsible for color vision in brighter light, and rods, which are essential for vision in dim light but do not perceive colors. After light passes through the cornea, lens system, and the vitreous humor, it enters the retina from the inside; that is, it passes through the
25 ganglion cells and nerve fibers, the inner and outer plexiform layers, the inner and outer nuclear layers, and the internal and external limiting membranes before it finally reaches the layer of photoreceptors located near the outside of the retina, just inside the outermost
30 pigment epithelium layer. The cells of the pigment epithelium layer act as an anatomical barrier to liquids and substances located outside of the eye, forming the "blood-retina" barrier, and provide nourishment, oxygen, a source of functionally useful substances like vitamin A,
35 and phagocytosis of decomposition products to photoreceptor

cells. There is no anatomical connection between the pigment epithelium and the photoreceptor layer, permitting separation of the layers in some pathological situations.

5 When rods or cones are excited by light, signals are transmitted through successive neurons in the retina itself, into the optic nerve fibers, and ultimately to the cerebral cortex. Both rods and cones contain molecules that decompose on exposure to light and, in the process, excite the nerve fibers leading from the eye. The molecule
10 in rods is rhodopsin. The three light-sensitive molecules in cones, collectively called iodopsin, have compositions only slightly different from that of rhodopsin and are maximally excited by red, blue, or green light, respectively.

15 Neither rods nor cones generate action potentials. Rather, the light-induced membrane hyperpolarization generated in the outer, photosensitive segment of a rod or cone cell is transmitted from the outer segment through the inner segment to the synaptic body by direct conduction of
20 the electrical voltage itself, a process called electrotonic conduction. At the synaptic body, the membrane potential controls the release of an unknown transmitter molecule. In low light, rod and cone cell membranes are depolarized and the rate of transmitter
25 release is greatest. Light-induced hyperpolarization causes a marked decrease in the release of transmitter molecules.

The transmitters released by rod and cone cells induce signals in the bipolar neurons and horizontal cells. The
30 signals in both these cells are also transmitted by electrotonic conduction and not by action potential.

The rod bipolar neurons connect with as many as 50 rod cells, while the dwarf and diffuse bipolar cells connect with one or several cone cells. A depolarizing bipolar
35 cell is stimulated when its connecting rods or cones are

exposed to light. The release of transmitter molecules inhibits the depolarizing bipolar cell. Therefore, in the dark, when the rods and cones are secreting large quantities of transmitter molecules, the depolarizing
5 bipolar cells are inhibited. In the light, the decrease in release of transmitter molecules from the rods and cones reduces the inhibition of the bipolar cell, allowing it to become excited. In this manner, both positive and negative signals can be transmitted through different bipolar cells
10 from the rods and cones to the amacrine and ganglion cells.

As their name suggests, horizontal cells project horizontally in the retina, where they may synapse with rods, cones, other horizontal cells, or a combination of cells types. The function of horizontal cells is unclear,
15 although some mechanism in the convergence of photoreceptor signaling has been postulated.

All types of bipolar cells connect with ganglion cells, which are of two primary types. A-type ganglion cells predominately connect with rod bipolar cells, while
20 B-type ganglion cells predominately connect with dwarf and diffuse bipolar cells. It appears that A-type ganglion cells are sensitive to contrast, light intensity, and perception of movement, while B-type ganglion cells appear more concerned with color vision and visual acuity.

25 Like horizontal cells, the Amacrine cells horizontally synapse with several to many other cells, in this case bipolar cells, ganglion cells, and other Amacrine cells. The function of Amacrine cells is also unclear.

The axons of ganglion cells carry signals into the
30 nerve fiber layer of the eye, where the axons converge into fibers which further converge at the optic disc, where they exit the eye as the optic nerve. The ganglion cells transmit their signals through the optic nerve fibers to the brain in the form of action potentials. These cells,
35 even when unstimulated, transmit continuous nerve impulses

at an average, baseline rate of about 5 per second. The visual signal is superimposed onto this baseline level of ganglion cell stimulation. It can be either an excitatory signal, with the number of impulses increasing above the baseline rate, or an inhibitory signal, with the number of nerve impulses decreasing below the baseline rate.

As part of the central nervous system, the eye is in some ways an extension of the brain; as such, it has a limited capacity for regeneration. This limited regeneration capacity further complicates the challenging task of improving vision, resolving dysfunction of the visual system, and/or treating or preventing ophthalmologic disorders. Many disorders of the eye, such as retinal photic injury, retinal ischemia-induced eye injury, age-related macular degeneration, free radical-induced eye diseases, as well as numerous other disorders, are considered to be entirely untreatable. Other ophthalmologic disorders, e.g., disorders causing permanent visual impairment, are corrected only by the use of ophthalmic devices and/or surgery, with varying degrees of success.

The immunosuppressant drugs FK506, rapamycin, and cyclosporin are well known as potent T-cell specific immunosuppressants, and are effective against autoimmunity, transplant or graft rejection, inflammation, allergic responses, other autoimmune or immune-mediated diseases, and infectious diseases. It has been disclosed that application of Cyclosporin, FK-506, Rapamycin, Buspirone, Spiperone, and/or their derivatives are effective in treating some ophthalmologic disorders of these types. Several ophthalmologic disorders or vision problems are known to be associated with autoimmune and immunologically-mediated activities; hence, immunomodulatory compounds are expected to demonstrate efficacy for treating those types of ophthalmologic disorders or vision problems.

The effects of FK506, Rapamycin, and related agents in the treatment of ophthalmologic diseases are disclosed in several U.S. patents (Goulet et al., U.S. Patent No. 5,532,248; Mochizuki et al., U.S. Patent No. 5,514,686; 5 Luly et al., U.S. Patent No. 5,457,111; Russo et al., U.S. Patent No. 5,441,937; Kulkarni, U.S. Patent No. 5,387,589; Asakura et al., U.S. Patent No. 5,368,865; Goulet et al., U.S. Patent No. 5,258,389; Armistead et al., U.S. Patent No. 5,192,773; Goulet et al., U.S. Patent No. 5,189,042; 10 and Fehr, U.S. Patent No. 5,011,844). These patents claim FK506 or Rapamycin related compounds and disclose the known use of FK506 or Rapamycin related compounds in the treatment of ophthalmologic disorders in association with the known immunosuppressive effects of FK506 and Rapamycin. 15 The compounds disclosed in these patents are relatively large. Further, the cited patents relate to immunomodulatory compounds limited to treating autoimmunity or related diseases, or immunologically-mediated diseases, for which the efficacy of FK506 and Rapamycin is well 20 known.

Other U.S. patents disclose the use of cyclosporin, Spiperone, Buspirone, their derivatives, and other immunosuppressive compounds for use in the treatment of ophthalmologic diseases (Sharpe et al., U.S. Patent No. 25 5,703,088; Sharpe et al., U.S. Patent No. 5,693,645; Sullivan, U.S. Patent No. 5,688,765; Sullivan, U.S. Patent No. 5,620,921; Sharpe et al., U.S. Patent No. 5,574,041; Eberle, U.S. Patent No. 5,284,826; Sharpe et al., U.S. Patent No. 5,244,902; Chiou et al., U.S. Patent Nos. 30 5,198,454 and 5,194,434; and Kaswan, U.S. Patent No. 4,839,342). These patents also relate to compounds useful for treating autoimmune diseases and cite the known use of cyclosporin, Spiperone, Buspirone, their derivatives, and other immunosuppressive compounds in treating ocular 35 inflammation and other immunologically-mediated

ophthalmologic diseases.

5 The immunosuppressive compounds disclosed in the prior art suppress the immune system, by definition, and also exhibit other toxic side effects. Accordingly, there is a need for non-immunosuppressant, small molecule compounds, and compositions and methods for use of such compounds, that are useful in improving vision; preventing, treating, and/or repairing visual impairment or dysfunction of the visual system; and preventing, treating, and/or resolving
10 ophthalmologic disorders.

There are also a number of patents on non-immunosuppressive compounds disclosing methods of use for permitting or promoting wound healing (whether from injury or surgery); controlling intraocular pressure (often
15 resulting from glaucoma); controlling neurodegenerative eye disorders, including damage or injury to retinal neurons, damage or injury to retinal ganglion cells, and macular degeneration; stimulating neurite outgrowth; preventing or reducing oxidative damage caused by free radicals; and
20 treating impaired oxygen and nutrient supply, as well as impaired waste product removal, resulting from low blood flow. These non-immunosuppressive substances fall into one of two general categories: naturally occurring molecules, such as proteins, glycoproteins, peptides, hormones, and
25 growth factors; and synthetic molecules.

Within the group of naturally occurring non-immunosuppressive molecules, several hormones, growth factors, and signaling molecules have been patented for use as supplements to naturally occurring quantities of such
30 molecules, as well as for targeting of specific cells where the particular molecule does not naturally occur in a mature individual. These patents generally claim methods of use for reducing or preventing the symptoms of ocular disease, or arresting or reversing vision loss.

35 Specifically, Louis et al., U.S. Patent Nos. 5,736,516

and 5,641,749, disclose the use of a glial cell line derived neurotrophic factor (GDNF) to stop or reverse the degeneration of retinal neurons (i.e. photoreceptors) and retinal ganglion cells caused by glaucoma, or other degenerative or traumatic retinal diseases or injuries. 5 O'Brien, et al., U.S. Patent Nos. 5,714,459 and 5,700,909, disclose the use of a glycoprotein, Saposin, and its derivatives for stimulating neurite outgrowth and increasing myelination. To stop or reverse degeneration 10 of retinal neurons, LaVail et al., U.S. Patent No. 5,667,968, discloses the use of a variety of neurotrophic proteins, including brain-derived neurotrophic factor, ciliary neurotrophic factor, neurotrophin-3 or neurotrophin-4, acidic or basic fibroblast growth factors, 15 interleukin, tumor necrosis factor- α , insulin-like growth factor-2 and other growth factors. Wong et al., U.S. Patent No. 5,632,984, discloses the use of interferons, especially interferon α -2a, for treating the symptoms of macular degeneration by reducing hemorrhage and limiting neovascularization. Finally, Wallace et al., U.S. Patent 20 No. 5,441,937, discloses the use of a lung-derived neurotrophic factor (NTF) to maintain the functionality of ciliary ganglion and parasympathetic neuron cells.

A key characteristic of factors derived from specific 25 cell lines is their localization to specific cell lines or tissues; systemic treatment with these molecules would run a substantial risk of unintended, and potentially dangerous, effects in cell lines where the genes encoding these molecules are inactive. Similarly, hormones and 30 growth factors often activate a large number of genes in many cell lines; again, non-localized application of these molecules would run a substantial risk of provoking an inappropriate, and potentially dangerous, response.

Within the category of synthetic molecules, most of 35 the patented compounds are immunosuppressive and disclose

uses in treating inflammatory, autoimmune, and allergic responses, as discussed above. A few others are non-immunosuppressive and claim the ability to treat cellular degeneration, and in some cases promote cellular regeneration, most often in the context of their antioxidant properties.

Specifically, Tso et al., U.S. Patent No. 5,527,533, discloses the use of astaxanthin, a carotenoid antioxidant, for preventing or reducing photoreceptor damage resulting from the presence of free radicals. Similarly, Babcock et al., U.S. Patent No. 5,252,319, discloses the use of antioxidant aminosteroids for treating eye disease and injury, by increasing resistance to oxidative damage. Freeman, U.S. Patent No. 5,468,752, discloses the use of the antiviral phosphonylmethoxyalkylcytosines to reduce abnormally increased intraocular pressure.

Naturally occurring hormones, growth factors, cytokines, and signaling molecules are generally multifunctional and activate many genes in diverse cell lines. The present compounds do not, thus avoiding the unexpected, and potentially dangerous, side effects of systemic use. Similarly, the present compounds also avoid the potential unexpected side effects of introducing cell line-specific molecules into other cell lines were they do not naturally occur.

The epithelial hair cells in the organ of Corti of the inner ear, transduce sound into neural activity, which is transmitted along the cochlear division of the eighth cranial nerve. This nerve consists of fibers from three types of neurons (Spoendlin, H. H., in Friedmann, I. Ballantyne, J., eds. "Ultrastructural Atlas of the Inner Ear", London, Butterworth, pp. 133-164, (1984)) 1) afferent neurons, which lie in the spiral ganglion and connect the cochlea to the brainstem; 2) efferent olivocochlear neurons, which originate in the superior olivary complex;

and 3) autonomic adrenergic neurons, which originate in the cervical sympathetic trunk and innervate the cochlea. In the human, there are approximately 30,000 afferent cochlear neurons, with myelinated axons, each consisting of about 50 lamellae, and 4-6 μ m in diameter. This histologic structure forms the basis of uniform conduction velocity, which is an important functional feature. Throughout the length of the auditory nerve, there is a trophic arrangement of afferent fibers, with 'basal' fibers wrapped over the centrally placed 'apical' fibers in a twisted rope-like fashion. Spoendlin (Spoendlin, H. H. in Naunton, R. F., Fernadex, C. eds., "Evoked Electrical Activity in the Auditory Nervous System", London, Academic Press, pp. 21-39, (1978)) identified two types of afferent neurons in the spiral ganglion on the basis of morphologic differences: type I cells (95%) are bipolar and have myelinated cell bodies and axons that project to the inner hair cells. Type II cells (5%) are monopolar with unmyelinated axons and project to the outer hair cells of the organ of Corti. Each inner hair cell is innervated by about 20 fibers, each of which synapses on only one cell. In contrast, each outer hair cell is innervated by approximately six fibers, and each fiber branches to supply approximately 10 cells. Within the cochlea, the fibers divide into: 1) an inner spiral group, which arises primarily ipsilaterally and synapses with the afferent neurons to the inner hair cells, and 2) a more numerous outer radial group, which arises mainly contralaterally and synapses directly with outer hair cells. There is a minimal threshold at one frequency, the characteristic or best frequency, but the threshold rises sharply for frequencies above and below this level (Pickles, J.O. in "Introduction to the Physiology of Hearing", London, Academic Press, pp. 71-106, (1982)). Single auditory nerve fibers therefore appear to behave as band-pass filters.

The basilar membrane vibrates preferentially to different frequencies, at different distances along its length, and the frequency selectivity of each cochlear nerve fiber is similar to that of the inner hair cell to which the fiber is connected. Thus, each cochlear nerve fiber exhibits a tuning curve covering a different range of frequencies from its neighboring fiber (Evans, E.F. in Beagley H.A. ed., "Auditory investigation: The Scientific and Technological basis", New York, Oxford University Pressm (1979)). By this mechanism, complex sounds are broken down into component frequencies (frequency resolution) by the filters of the inner ear.

Impairment anywhere along the auditory pathway, from the external auditory canal to the central nervous system, may result in hearing loss. The auditory apparatus can be subdivided into the external and middle ear, inner ear and auditory nerve and central auditory pathways. Auditory information in humans is transduced from a mechanical signal to a neurally conducted electrical impulse by the action of approximately 15,000 epithelial cells (hair cells) and 30,000 first-order neurons (spiral ganglion cells) in the inner ear. All central fibers of spiral ganglion neurons form synapses in the cochlear nucleus of the pontine brainstem. The number of neurons involved in hearing increases dramatically from the cochlea to the auditory brain stem and the auditory cortex. All auditory information is transduced by only 15,000 hair cells, of which the so-called inner hair cells, numbering 3500, are critically important, since they form synapses with approximately 90 percent of the 30,000 primary auditory neurons. Thus, damage to a relatively few cells in the auditory periphery can lead to substantial hearing loss. Hence, most causes of sensorineural loss can be ascribed to lesions in the inner ear (Nadol, J.B., New England Journal of Medicine, (1993), 329:1092-1102).

Hearing loss can be on the level of conductivity, sensorineural and central level. Conductive hearing loss is caused by lesions involving the external or middle ear, resulting in the destruction of the normal pathway of airborne sound amplified by the tympanic membrane and the ossicles to the inner ear fluids. Sensorineural hearing loss is due to lesions of the central auditory pathways. These consist of the cochlear and dorsal olivary nucleus complex, inferior colliculi, medial geniculate bodies, auditory cortex in the temporal lobes and interconnecting afferent and efferent fiber tracts (Adams R.D. and Maurice, V., eds., in "Principles of Neurology", (1989), McGraw-Hill Information Services Company, pp. 226-246).

Trauma due to acoustic overstimulation is another leading cause of deafness. There is individual susceptibility to trauma from noise. Clinically important sensorineural hearing loss may occur in some people exposed to high-intensity noise, even below levels approved by the Occupational Safety and Health Agency (Osguthorpe, J.D., ed., Washington D.C., American Academy of Otolaryngology-Head and Neck Surgery Foundation, (1988)).

Demyelinating processes, such as multiple sclerosis, may cause sensorineural hearing loss (Noffsinger, D., et al., Acto Otolaryngol. Suppl. (Stockh.) (1972), 303:1-63). More recently, a form of immune-mediated sensorineural hearing loss has been recognized (McCabe, B.F., Ann. Otol. Rhinol. Laryngol. (1979), 88:585-9). The hearing loss is usually bilateral, is rapidly progressive (measured in weeks and months), and may or may not be associated with vestibular symptoms.

A variety of tumors, both primary and metastatic, can produce either a conductive hearing loss, or a sensorineural hearing loss, by invading the inner ear or auditory nerve (Houck, J.R., et al., Otolaryngol. Head Neck Surg. (1992), 106:92-7). A variety of degenerative

disorders of unknown cause can produce sensorineural hearing loss. Meniere's syndrome (Nadol, J.B., ed., "Meniere's Disease: Pathogenesis, Pathophysiology, Diagnosis, And Treatment," Amsterdam: Kugler & Ghedini
5 (1989)), characterized by fluctuating sensorineural hearing loss, episodic vertigo, and tinnitus, appears to be caused by a disorder of fluid homeostasis within the inner ear, although the pathogenesis remains unknown. Sudden idiopathic sensorineural hearing loss (Wilson, W.R., et
10 al., Arch. Otolaryngol. (1980), 106:772-6), causing moderate-to-severe sensorineural deafness, may be due to various causes, including inner ear ischemia and viral labyrinthitis.

Regardless of the cause, there exists a need to
15 prevent or treat sensorineural hearing loss. The present invention provides such a method.


BRIEF SUMMARY OF THE INVENTION

The present invention is directed to novel
hydantoin derivative compounds, and their preparation
5 and use for treating neurodegenerative disorders, for
treating alopecia and related hair loss disorders, for
treating vision disorders and/or improving vision, for
treating memory impairment and/or enhancing memory
performance, and for treating sensorineural hearing
10 loss.

These compounds stimulate neuronal regeneration and
outgrowth and as such are useful for treating
neurological disorders and neurodegenerative diseases.
These compounds also promote hair growth and as such are
15 useful for treating hair loss disorders. These
compounds also are useful for treating vision disorders,
improving vision, treating memory impairment, enhancing
memory performance, or treating hearing loss. A
preferred feature of the compounds of the present
20 invention is that they do not exert any significant
immunosuppressive activity and/or are non-
immunosuppressive.

A preferred embodiment of this invention is a
compound having the formula I:

25



I

where

each X independently is O, S, or NR₂;
30 R₂ is selected from the group consisting of
cyano, nitro, hydrogen, C₁-C₄ alkyl, hydroxy,
and C₁-C₄ alkoxy;

D is a direct bond or a C₁-C₈ alkyl or alkenyl;

R is selected from the group consisting of
hydrogen, phenyl, biphenyl, cyclopropyl,
cyclobutyl, cyclopentyl, cycloheptyl,
5 cyclooctyl, naphthyl, 1,2,3,4-
tetrahydronaphthyl, indenyl, azulenyl,
fluorenyl, anthracenyl, isoindolyl, indoliny,
benzofuranyl, benzothiophenyl, indazolyl,
benzimidazolyl, tetrahydrofuranyl,
10 tetrahydropyranyl, pyridyl, pyrrolyl,
pyrrolidinyl, pyridinyl, pyrimidinyl, purinyl,
quinolinyl, isoquinolinyl,
tetrahydroquinolinyl, quinoliziny, furyl,
benzofuranyl, thiophenyl, imidazolyl,
15 oxazolyl, benzoxazolyl, benzoxazinyl,
thiazolyl, isoxazolyl, isotriazolyl,
oxadiazolyl, triazolyl, thiadiazolyl,
pyridazinyl, pyrimidinyl, pyrazinyl,
triazinyl, trithianyl, indoliziny, pyrazolyl,
20 pyrazolinyl, pyrazolidinyl, benzopyranyl,
thienyl, tetrahydroisoquinolinyl, cinnolinyl,
phthalazinyl, quinazolinyl, quinoxalinyl,
naphthyridinyl, pteridinyl, carbazolyl,
phenazinyl, phenothiazinyl, phenoxazinyl, and
25 adamantyl;

wherein R may be optionally substituted with one
substituent which is selected from the group
consisting of hydrogen, halo, hydroxyl, nitro,
trifluoromethyl, C₁-C₆ straight or branched chain
30 alkyl, C₂-C₆ straight or branched chain alkenyl, C₁-
C₄ alkoxy, C₂-C₄ alkenyloxy, phenyl, phenoxy,
benzyloxy, and amino;

or a pharmaceutically acceptable salt, ester, or solvate
thereof;


35 wherein when R is hydrogen, D is a C₅-C₇ alkyl or C₅-

C₈ alkenyl;

wherein when R is phenyl, R must be substituted with phenyl, hydroxyl, trifluoromethyl, C₂-C₆ straight or branched chain alkyl or alkenyl, C₃-C₄ alkoxy or C₂-C₄ alkenyloxy, phenoxy, or benzyloxy;

wherein when R is 4-trifluoromethylphenyl, both X substituents must be O.

Another preferred embodiment of this invention is a compound of formula II:



II

where

each X independently is O, S, or NR₂;

R₂ is selected from the group consisting of cyano, nitro, hydrogen, C₁-C₄ alkyl, hydroxy, and C₁-C₄ alkoxy;

D is a direct bond or a C₁-C₈ alkyl or alkenyl;

R is selected from the group consisting of

hydrogen, phenyl, biphenyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, 1,2,3,4-tetrahydronaphthyl, indenyl, azulenyl, fluorenyl, anthracenyl, indolyl, isoindolyl, indolinyl, benzofuranyl, benzothiophenyl, indazolyl, benzimidazolyl, tetrahydrofuranyl, tetrahydropyranyl, pyridyl, pyrrolyl, pyrrolidinyl, pyridinyl, pyrimidinyl, purinyl, quinolinyl, isoquinolinyl, tetrahydroquinolinyl, quinolizinyl, furyl, thiophenyl, imidazolyl, oxazolyl,

benzopyranyl, thiazolyl, isotriazolyl,
oxadiazolyl, triazolyl, thiadiazolyl,
pyridazinyl, pyrimidinyl, pyrazinyl,
triazinyl, trithianyl, indolizinyl, pyrazolyl,
5 pyrazolinyl, pyrazolidinyl, thienyl,
tetrahydroisoquinolinyl, cinnolinyl,
phthalazinyl, quinazolinyl, naphthyridinyl,
pteridinyl, carbazolyl, acridinyl, phenazinyl,
phenothiazinyl, phenoxazinyl, and adamantyl;

10 wherein R may be optionally substituted with one
substituent which is selected from the group
consisting of hydrogen, halo, hydroxyl, nitro,
trifluoromethyl, C₁-C₆ straight or branched chain
alkyl, C₂-C₆ straight or branched chain alkenyl, C₁-
15 C₄ alkoxy, C₂-C₄ alkenyloxy, phenyl, phenoxy,
benzyloxy, and amino;


or a pharmaceutically acceptable salt, ester, or solvate
thereof;

20 wherein when R is hydrogen, D is a C₅-C₈ alkyl or
alkenyl;

wherein when R is phenyl, R must be substituted
with C₂-C₃ or C₅-C₆ straight or branched chain alkyl or
alkenyl, C₃-C₄ alkoxy or C₂-C₄ alkenyloxy, phenyl,
phenoxy, benzyloxy, or amino.

25 Another preferred embodiment of this invention is a
pharmaceutical composition containing:

(i) an effective amount of a hydantoin derivative
compound of formula III:



III

30 where

each X independently is O, S, or NR₂;

R₂ is selected from the group consisting of

cyano, nitro, hydrogen, C₁-C₄ alkyl, hydroxy,
and C₁-C₄ alkoxy;

D is a direct bond or C₁-C₈ alkyl or alkenyl;


R is hydrogen, or an alicyclic or aromatic,
mono-, bi- or tricyclic, carbo- or
heterocyclic ring;

wherein R is optionally substituted with one
substituent selected from the group consisting of
hydrogen, halo, hydroxyl, nitro, trifluoromethyl,
C₁-C₆ straight or branched chain alkyl, C₂-C₆
straight or branched chain alkenyl, C₁-C₄ alkoxy, C₂-
C₄ alkenyloxy, phenyl, phenoxy, benzyloxy, and
amino; and

(ii) a pharmaceutically acceptable carrier.

Another preferred embodiment of this invention is a
pharmaceutical composition containing:

(i) an effective amount of a hydantoin derivative
compound of formula IV:



IV

where

each X independently is O, S, or NR₂;

R₂ is selected from the group consisting of
cyano, nitro, hydrogen, C₁-C₄ alkyl, hydroxy,
and C₁-C₄ alkoxy;

D is a direct bond or C₁-C₈ alkyl or alkenyl;

R is hydrogen, or an alicyclic or aromatic,
mono-, bi- or tricyclic, carbo- or
heterocyclic ring;

wherein R is optionally substituted with one
substituent selected from the group consisting of
hydrogen, halo, hydroxyl, nitro, trifluoromethyl,

C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or branched chain alkenyl, C₁-C₄ alkoxy, C₂-C₄ alkenyloxy, phenyl, phenoxy, benzyloxy, and amino; and

5 (ii) a pharmaceutically acceptable carrier.

For neurotrophic compositions, a neurotrophic factor different from the present inventive compounds may also be administered or otherwise included in the composition.

10 Another preferred embodiment of the invention is a method of treating a neurological disorder in an animal, comprising administering to the animal an effective amount of the compound of formulae III or IV to stimulate growth of damaged peripheral nerves or to
15 promote neuronal regeneration.

Another preferred embodiment of the invention is a method of stimulating growth of damaged peripheral nerves, comprising administering to a damaged peripheral nerve an effective amount of the compound of formulae
20 III or IV to stimulate or promote growth of the damaged peripheral nerve.

Another preferred embodiment of the invention is a method of promoting neuronal regeneration and growth in animals, comprising administering to an animal an
25 effective amount of the compound of formulae III or IV to promote neuronal regeneration.

Another preferred embodiment of the invention is a method of preventing neurodegeneration in animals, comprising administering to an animal an effective
30 amount of the compound of formulae III or IV to prevent neurodegeneration.

Another preferred embodiment of the invention is a method for treating alopecia or promoting hair growth in an animal, comprising administering to an animal an
35 effective amount of the compound of formulae III or IV.

Another preferred embodiment of this invention is a method for treating a vision disorder, improving vision, treating memory impairment, enhancing memory performance, or treating sensorineural hearing loss in
5 an animal, comprising administering to an animal an effective amount of the compound of formula III or IV.

The present invention further contemplates a process for preparing the hydantoin derivative compounds of the invention, comprising acidifying an intermediate
10 compound.

The present invention further contemplates the compound(s) of the invention for use in treatment of a disease. In particular, the present invention contemplates the compound(s) of the invention for use in
15 treatment of the disorders enumerated herein.

The invention further contemplates the compound(s) of the invention for use in the preparation of a medicament or pharmaceutical composition. In particular, the invention contemplates the compound(s)
20 of the invention for use in the preparation of a medicament or pharmaceutical composition for treatment of the disorders enumerated herein.

The invention also provides for the use of compound(s) of the invention for treating a disease. In particular, the invention provides for the use of
25 compound(s) of the invention for treating the disorders enumerated herein.

The invention also provides for the use of compound(s) of the invention in the manufacture of a
30 medicament or pharmaceutical composition. In particular, the invention provides for the use of compound(s) of the invention in the manufacture of a medicament or pharmaceutical composition for the treatment of the disorders enumerated herein. Such
35 pharmaceutical compositions include, as appropriate to

the specific disorder, topical, systemic, oral or injectable formulations. It is further contemplated that the compound(s) of the invention may be administered with an effective amount of a second
5 therapeutic agent for the treatment of the enumerated disorders. A variety of pharmaceutical formulations and different delivery techniques are described in further detail below.

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
2218
2219
2220
2221
2222
2223
2224
2225
2226
22

DETAILED DESCRIPTION OF THE INVENTION

Definitions

"Alkenyl" means a branched or unbranched unsaturated hydrocarbon chain comprising a designated number of carbon atoms. For example, C₂-C₆ straight or branched alkenyl hydrocarbon chain contains 2 to 6 carbon atoms having at least one double bond, and includes but is not limited to substituents such as ethenyl, propenyl, iso-propenyl, butenyl, iso-butenyl, tert-butenyl, n-pentenyl, n-hexenyl, and the like. It is also contemplated as within the scope of the present invention that "alkenyl" may also refer to an unsaturated hydrocarbon chain wherein any of the carbon atoms of said alkenyl are optionally replaced with O, NH, S, or SO₂. For example, carbon 2 of 4-pentene can be replaced with O to form (2-propene)oxymethyl.

"Alkoxy" means the group -OR wherein R is alkyl as herein defined. Preferably, R is a branched or unbranched saturated hydrocarbon chain containing 1 to 6 carbon atoms.

"Alkyl" means a branched or unbranched saturated hydrocarbon chain comprising a designated number of carbon atoms. For example, C₁-C₆ straight or branched alkyl hydrocarbon chain contains 1 to 6 carbon atoms, and includes but is not limited to substituents such as methyl, ethyl, propyl, iso-propyl, butyl, iso-butyl, tert-butyl, n-pentyl, n-hexyl, and the like. It is also contemplated as within the scope of the present invention that "alkyl" may also refer to a hydrocarbon chain wherein any of the carbon atoms of said alkyl are optionally replaced with O, NH, S, or SO₂. For example, carbon 2 of n-pentyl can be replaced with O to form propyloxymethyl.

"Alopecia" refers to deficient hair growth and


partial or complete loss of hair, including without limitation androgenic alopecia (male pattern baldness), toxic alopecia, alopecia senilis, alopecia areata, alopecia pelada and trichotillomania. Alopecia results when the pilar cycle is disturbed. The most frequent phenomenon is a shortening of the hair growth or anagen phase due to cessation of cell proliferation. This results in an early onset of the catagen phase, and consequently a large number of hairs in the telogen phase during which the follicles are detached from the dermal papillae, and the hairs fall out. Alopecia has a number of etiologies, including genetic factors, aging, local and systemic diseases, febrile conditions, mental stresses, hormonal problems, and secondary effects of drugs.

"Aralkyl" refers to alkyl or alkylene (alkenyl) chain which is substituted with aryl, heteroaryl, carbocycle or heterocycle, or alternatively one or more aryl, heteroaryl, carbocycle, or heterocycle(s) which is/are substituted with alkyl or alkenyl, i.e. 'Alkyl/alkylene which is substituted with Ar' or 'Ar which is substituted with alkyl/alkylene'.

"Aryl" or "aromatic" refers to an aromatic carbocyclic or heterocyclic group having a single ring, for example a phenyl ring; multiple rings, for example biphenyl; or multiple condensed rings in which at least one ring is aromatic, for example naphthyl, 1,2,3,4-tetrahydronaphthyl, anthryl, or phenanthryl, which can be unsubstituted or substituted with one or more other substituents as defined above. The substituents attached to a phenyl ring portion of an aryl moiety in the compounds of the invention may be configured in the ortho-, meta-, or para- orientations orientations.

Examples of typical aryl moieties included in the scope of the present invention may include, but are not

limited to, the following:



5 "Carbocycle" or "carbocyclic" refers to an organic cyclic moiety in which the cyclic skeleton is comprised of only carbon atoms, whereas the term "heterocycle" or "heterocyclic" refers to an organic cyclic moiety in which the cyclic skeleton contains one or more heteroatoms selected from nitrogen, oxygen, or sulfur, and which may or may not include carbon atoms. Thus, the term "carbocycle" refers to a carbocyclic moiety containing the indicated number of carbon atoms. The term "C₃-C₈ cycloalkyl", therefore, refers to an organic cyclic substituent in which three to eight carbon atoms form a
10 three, four, five, six, seven, or eight-membered ring, including, for example, a cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, or cyclooctyl ring.

As used herein, "carbocyclic" or "heterocyclic" may also refer to two or more cyclic ring systems, which are
20 fused to form, for example, bicyclic, tricyclic, or other similar bridged ring substituents (e.g. adamantyl) or

multiple condensed ring systems.

"Enhancing memory performance" refers to improving or increasing the mental faculty by which to register, retain or recall past experiences, knowledge, ideas, sensations, thoughts or impressions.

"Eye" refers to the anatomical structure responsible for vision in humans and other animals, and encompasses the following anatomical structures, without limitation: lens, vitreous body, ciliary body, posterior chamber, anterior chamber, pupil, cornea, iris, canal of Schlemm, zonules of Zinn, limbus, conjunctiva, choroid, retina, central vessels of the retina, optic nerve, fovea centralis, macula lutea, and sclera.

"Halo" means at least one fluoro, chloro, bromo, or iodo moiety.


"Heterocycle" or "heterocyclic", as used herein, refers to a saturated, unsaturated or aromatic carbocyclic group having a single ring, multiple fused rings (for example, bicyclic, tricyclic, or other similar bridged ring systems or substituents), or multiple condensed rings, and having at least one hetero atom such as nitrogen, oxygen or sulfur within at least one of the rings. This term also includes "Heteroaryl" which refers to a heterocycle in which at least one ring is aromatic. Any heterocyclic or heteroaryl group can be unsubstituted or optionally substituted with one or more groups, as defined above. Further, bi- or tricyclic heteroaryl moieties may comprise at least one ring which is either completely or partially saturated.

In the context of the invention, useful carbo- and heterocyclic rings include, for example and without limitation, phenyl, benzyl, naphthyl, indenyl, azulenyl, fluorenyl, anthracenyl, indolyl, isoindolyl, indolinyl, benzofuranyl, benzothiophenyl, indazolyl, benzimidazolyl, benzthiazolyl, tetrahydrofuranyl, tetrahydropyranyl,


pyridyl, pyrrolyl, pyrrolidinyl, pyridinyl, pyrimidinyl,
purinyl, quinolinyl, isoquinolinyl, tetrahydroquinolinyl,
quinolizinyl, furyl, thiophenyl, imidazolyl, oxazolyl,
5 benzoxazolyl, thiazolyl, isoxazolyl, isotriazolyl,
oxadiazolyl, triazolyl, thiadiazolyl, pyridazinyl,
pyrimidinyl, pyrazinyl, triazinyl, trithianyl, indolizinyl,
pyrazolyl, pyrazolinyl, pyrazolidinyl, thienyl,
tetrahydroisoquinolinyl, cinnolinyl, phthalazinyl,
quinazolinyl, quinoxalinyl, naphthyridinyl, pteridinyl,
10 carbazolyl, acridinyl, phenazinyl, phenothiazinyl,
phenoxazinyl, and adamantyl.

As one skilled in the art will appreciate, such
heterocyclic moieties may exist in several isomeric forms,
all of which are encompassed by the present invention. For
15 example, a 1,3,5-triazine moiety is isomeric to a 1,2,4-
triazine group. Such positional isomers are to be
considered within the scope of the present invention.
Likewise, the heterocyclic or heteroaryl groups can be
bonded to other moieties in the compounds of the present
20 invention. The point(s) of attachment to these other
moieties is not to be construed as limiting on the scope
of the invention. Thus, by way of example, a pyridyl
moiety may be bound to other groups through the 2-, 3-, or
4-position of the pyridyl group. All such configurations
25 are to be construed as within the scope of the present
invention.

Examples of heterocyclic or heteroaryl moieties
included in the scope of the present invention may include,
but are not limited to, the following:




The Condensed Chemical Dictionary (Tenth Edition)
 revised by Gessner G. Hawley (1981) defines "hydantoins"
 as compounds having the chemical formula NHCONHCOCH_2 . The
 "hydantoin derivatives" contemplated by the present
 invention have the general structure:




wherein the five membered ring depicted is part of a
 bicyclic ring structure and wherein each X independently
 can be O, S, or NR_2 . In particular, one class of hydantoin

derivative compounds contemplated by the present invention has the structure:



5 Another class of hydantoin derivative compounds contemplated by the present invention has the structure:



"Isomers" are different compounds that have the same
10 molecular formula and includes cyclic isomers such as (iso)indole and other isomeric forms of cyclic moieties. "Stereoisomers" are isomers that differ only in the way the atoms are arranged in space. "Enantiomers" are a pair of stereoisomers that are non-superimposable mirror images of
15 each other. "Diastereoisomers" are stereoisomers which are not mirror images of each other. "Racemic mixture" means a mixture containing equal parts of individual enantiomers. "Non-racemic mixture" is a mixture containing unequal parts of individual enantiomers or stereoisomers.

20 "Memory impairment" refers to a diminished mental registration, retention or recall of past experiences, knowledge, ideas, sensations, thoughts or impressions. Memory impairment may affect short and long-term information retention, facility with spatial relationships,
25 memory (rehearsal) strategies, and verbal retrieval and production. Common causes of memory impairment are age, severe head trauma, brain anoxia or ischemia, alcoholic-nutritional diseases, and drug intoxications. Examples of memory impairment include, without limitation, benign

forgetfulness, amnesia and any disorder in which memory deficiency is present, such as Korsakoff's amnesic psychosis, dementia and learning disorders.

5 The term "middle-ear" refers to the space between the tympanic membrane and the inner ear. This location is external to all inner ear tissue and an invasive procedure might not be required to access this region if a formulation capable of penetrating through the tympanic membrane is administered. Otherwise, the material will be
10 introduced to the middle ear by injection through the tympanic membrane or, in case repeated administrations are needed, a hole can be made in the tympanic membrane. An opening in the tympanic membrane is a frequent procedure, performed on an office-visit basis, in cases such as
15 infections of the middle ear (usually in children). The opening generally closes spontaneously after a few days.

"Neopsic factors" or "neopsics" refers to compounds useful in treating vision loss, preventing vision degeneration, or promoting vision regeneration.

20 "Neopsis" refers to the process of treating vision loss, preventing vision degeneration, or promoting vision regeneration.

"Neurotrophic" as used herein includes without limitation the ability to stimulate neuronal regeneration or growth, and/or the ability to prevent or treat
25 neurodegeneration.

"Non-immunosuppressive" as used herein refers to the inability of the compounds of the present invention to trigger an immune response when compared to a control such
30 as FK506 or cyclosporin A. Assays for determining immunosuppression are well known to those of ordinary skill in the art. Specific, non-limiting examples of well known assays include PMA and OKT3 wherein mitogens are used to stimulate proliferation of human peripheral blood
35 lymphocytes (PBC) and the compounds are evaluated on their

ability to inhibit such proliferation.

"Ophthalmological" refers to anything about or concerning the eye, without limitation, and is used interchangeably with "ocular," "ophthalmic,"
5 "ophthalmologic," and other such terms, without limitation.

"Pharmaceutically acceptable carrier" as used herein refers to any carrier, diluent, excipient, suspending agent, lubricating agent, adjuvant, vehicle, delivery system, emulsifier, disintegrant, absorbant, preservative,
10 surfactant, colorant, flavorant, or sweetener. For these purposes, the compounds of the present invention may be administered orally, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally or via an implanted reservoir in dosage formulations containing
15 conventional non-toxic pharmaceutically-acceptable carriers, adjuvants and vehicles. The term parenteral as used herein includes subcutaneous, intravenous, intramuscular, intraperitoneally, intrathecally, intraventricularly, intrasternal and intracranial injection or infusion techniques.
20

"Pharmaceutically acceptable salt", as used herein, refers to an organic or inorganic salt which is useful in the treatment of a warm-blooded animal in need thereof. Such salts can be acid or basic addition salts, depending
25 on the nature of the inventive compound to be used.

Further, the compounds of the invention may exist as crystalline solids which can be crystallized from common solvents such as ethanol, N,N-dimethyl-formamide, water, or the like. Thus, crystalline forms of the compounds of
30 the invention may exist as solvates and/or hydrates of the parent compounds or their pharmaceutically acceptable salts. All of such forms likewise are to be construed as falling within the scope of the invention.

"Pilar cycle" refers to the life cycle of hair
35 follicles, and includes three phases:

- (1) the anagen phase, the period of active hair growth which, insofar as scalp hair is concerned, lasts about three to five years;
- 5 (2) the catagen phase, the period when growth stops and the follicle atrophies which, insofar as scalp hair is concerned, lasts about one to two weeks; and
- 10 (3) the telogen phase, the rest period when hair progressively separates and finally falls out which, insofar as scalp hair is concerned, lasts about three to four months.

Normally 80 to 90 percent of the follicles are in the anagen phase, less than 1 percent being in the catagen phase, and the rest being in the telogen phase. In the
15 telogen phase, hair is uniform in diameter with a slightly bulbous, non-pigmented root. By contrast, in the anagen phase, hair has a large colored bulb at its root.

"Preventing neurodegeneration" as used herein includes the ability to inhibit or prevent neurodegeneration in
20 patients newly diagnosed as having a neurodegenerative disease, or at risk of developing a new degenerative disease and for inhibiting or preventing further neurodegeneration in patients who are already suffering from or have symptoms of a neurodegenerative disease when
25 the compounds are given concurrently.

"Preventing vision degeneration" as used herein includes the ability to prevent degeneration of vision in patients newly diagnosed as having a degenerative disease affecting vision, or at risk of developing a new
30 degenerative disease affecting vision, and for preventing further degeneration of vision in patients who are already suffering from or have symptoms of a degenerative disease affecting vision.

"Primary ring structure" refers to a 5- or 6-membered
35 ring structure which is depicted in the formula drawings

herein. Such definition shall apply to only one ring structure in any molecule described in this application, regardless of the number or confirmation of any substituent cyclic structures.

5 "Promoting hair growth" refers to maintaining, inducing, stimulating, accelerating, or revitalizing the germination of hair.

"Promoting vision regeneration" refers to maintaining, improving, stimulating or accelerating recovery of, or
10 revitalizing one or more components of the visual system in a manner which improves or enhances vision, either in the presence or absence of any ophthalmologic disorder, disease, or injury.

The term "small molecule" refers to the molecular
15 weight of the compounds of the invention as compared to FK506. Thus, the term "small molecule" includes molecular weights less than about 800 Daltons (m.w.), and novel subranges or limits below the same including about 100 to about 750 m.w., about 150 to about 500 m.w., about 150 to
20 about 350 m.w., about 200 to about 300 m.w., about 210 to about 280 m.w., about 220 to about 260, and about 240 m.w.. The term "spatially small molecule" refers to the capability of the compounds to fit entirely or substantially within the binding cavity of FKBP-12 as
25 compared to FK506.

"Treating" or "treatment" as used herein covers any treatment of a disease and/or condition in an animal, particularly a human, and includes:

(i) preventing a disease and/or condition from
30 occurring in a subject which may be predisposed to the disease and/or condition but has not yet been diagnosed as having it;

(ii) inhibiting the disease and/or condition, i.e., arresting its development; or

35 (iii) relieving the disease and/or condition, i.e.,

causing regression of the disease and/or condition.

"Treating alopecia" refers to:

(i) preventing alopecia in an animal which may be predisposed to alopecia; and/or

5 (ii) inhibiting, retarding or reducing alopecia; and/or

(iii) promoting hair growth; and/or

(iv) prolonging the anagen phase of the hair cycle; and/or

10 (v) converting vellus hair to growth as terminal hair.

Terminal hair is coarse, pigmented, long hair in which the bulb of the hair follicle is seated deep in the dermis. Vellus hair, on the other hand, is fine, thin, non-pigmented short hair in which the hair bulb is located
15 superficially in the dermis. As alopecia progresses, the hairs change from the terminal to the vellus type.

"Vision", as used herein, refers to the ability of humans and other animals to process images, and is used interchangeably with "sight", "seeing", and other such
20 terms, without limitation.


"Vision disorder" refers to any disorder that affects or involves vision, including without limitation visual impairment, orbital disorders, disorders of the lacrimal apparatus, disorders of the eyelids, disorders of the
25 conjunctiva, disorders of the cornea, cataracts, disorders of the uveal tract, disorders of the optic nerve or visual pathways, free radical induced eye disorders and diseases, immunologically-mediated eye disorders and diseases, eye injuries, and symptoms and complications of eye disease,
30 eye disorder, or eye injury.

"Visual impairment" refers to any dysfunction in vision including without limitation disturbances or diminution in vision (e.g., binocular, central, peripheral, scotopic), visual acuity for objects near and for, visual
35 field, ocular motility, color perception, adaptation to

light and dark, accommodation, refraction, and lacrimation.
See *Physicians' Desk Reference (PDR) for Ophthalmology*,
16th Edition, 6:47 (1988).

5 "Warm-blooded animal" includes a mammal, including a
member of the human, equine, porcine, bovine, murine,
canine or feline species. In the case of a human, the term
"warm-blooded animal" may also be referred to as a
"patient". Further, as used herein, "a warm blooded animal
10 in need thereof" refers to a warm-blooded animal which is
susceptible to a disorder due to genetic or environmental
conditions or predispositions. This term also refers to
a warm blooded animal which has already suffered some
degree of injury or damage because of genetic or
15 environmental conditions to which the animal has been
exposed or to which it has been predisposed. Environmental
conditions can include treatment with a therapeutic
compound, as well as other types of injury or insult.

Further, as used throughout the teaching of the
invention, a designation of:



wherein W or Y is H₂, or similar designations, is meant to
25 denote that two hydrogen atoms are attached to the noted
carbon and that the bonds to each hydrogen are single
bonds.

Compounds of the Invention

The present invention relates to the surprising discovery that the inventive hydantoin and hydantoin derivative compounds are neurotrophic, are able to treat alopecia, are able to treat vision and memory disorders, and are able to treat sensorineural hearing loss. Accordingly, a novel class of hydantoin derivative compounds is provided. A preferred feature of the compounds of the present invention is that they do not exert any significant immunosuppressive activity.

The neurotrophic compounds of this invention can be periodically administered to a patient undergoing treatment for neurological disorders or for other reasons in which it is desirable to stimulate neuronal regeneration and growth, such as in various peripheral neuropathic and neurological disorders relating to neurodegeneration. The compounds of this invention can also be administered to mammals other than humans for treatment of various mammalian neurological disorders.

The novel compounds of the present invention possess an excellent degree of neurotrophic activity. This activity is useful in the stimulation of damaged neurons, the promotion of neuronal regeneration, the prevention of neurodegeneration, and in the treatment of several neurological disorders known to be associated with neuronal degeneration and peripheral neuropathies. The neurological disorders that may be treated include but are not limited to: trigeminal neuralgia, glossopharyngeal neuralgia, Bell's Palsy, myasthenia gravis, muscular dystrophy, amyotrophic lateral sclerosis, progressive muscular atrophy, progressive bulbar inherited muscular atrophy, herniated, ruptured or prolapsed invertebrate disk syndromes, cervical spondylosis, plexus disorders, thoracic outlet

destruction syndromes, peripheral neuropathic such as those caused by lead, dapsone, ticks, prophyria, or Gullain-Barré syndrome, multiple sclerosis, stroke and ischemia associated with stroke, neural paropathy, other neurodegenerative diseases, motor neuron diseases, sciatic crush, peripheral neuropathy, particularly neuropathy associated with diabetes, spinal cord injuries and facial nerve crush, Huntington's Disease, Alzheimer's disease, and Parkinson's disease.

10 The above discussion relating to the utility and administration of the compounds of the present invention also applies to the pharmaceutical compositions of the present invention.

15 For these purposes, the compounds of the present invention may be administered orally, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally or via an implanted reservoir in dosage formulations containing conventional non-toxic pharmaceutically-acceptable carriers, adjuvants and vehicles. The term parenteral as used herein includes subcutaneous, intravenous, intramuscular, intraperitoneally, intrathecally, intraventricularly, intrasternal and intracranial injection or infusion techniques.

25 For oral administration, the compounds of the present invention may be provided in any suitable dosage form known in the art. For example, the compositions may be incorporated into tablets, powders, granules, beads, chewable lozenges, capsules, liquids, aqueous suspensions or solutions, or similar dosage forms, using conventional equipment and techniques known in the art. Tablet dosage forms are preferred. Tablets may contain carriers such as lactose and corn starch, and/or lubricating agents such as magnesium stearate. Capsules may contain diluents including lactose and dried corn

starch. Aqueous suspensions may contain emulsifying and suspending agents combined with the active ingredient.

When preparing dosage form incorporating the compositions of the invention, the compounds may also be
5 blended with conventional excipients such as binders, including gelatin, pregelatinized starch, and the like; lubricants, such as hydrogenated vegetable oil, stearic acid, and the like; diluents, such as lactose, mannose, and sucrose; disintegrants, such as
10 carboxymethylcellulose and sodium starch glycolate; suspending agents, such as povidone, polyvinyl alcohol, and the like; absorbants, such as silicon dioxide; preservatives, such as methylparaben, propylparaben, and sodium benzoate; surfactants, such as sodium lauryl
15 sulfate, polysorbate 80, and the like; colorants such as F.D. & C. dyes and lakes; flavorants; and sweeteners.

Compositions and methods of the invention also may utilize controlled release technology. Thus, for example, the inventive compounds may be incorporated
20 into a hydrophobic polymer matrix for controlled release over a period of days. Such controlled release films are well known to the art. Particularly preferred are transdermal delivery systems. Other examples of polymers commonly employed for this purpose that may be
25 used in the present invention include nondegradable ethylene-vinyl acetate copolymer and degradable lactic acid-glycolic acid copolymers which may be used externally or internally. Certain hydrogels such as poly(hydroxyethylmethacrylate) or poly(vinylalcohol)
30 also may be useful, but for shorter release cycles than the other polymer releases systems, such as those mentioned above.

To be effective therapeutically as central nervous system targets, the compounds of the present invention
35 should readily penetrate the blood-brain barrier when

peripherally administered. Compounds which cannot
penetrate the blood-brain barrier can be effectively
administered by an intraventricular route or other
appropriate delivery system suitable for administration
5 to the brain.

The compounds of the present invention may be
administered in the form of sterile injectable
preparations, for example, as sterile injectable aqueous
or oleaginous suspensions. These suspensions may be
10 formulated according to techniques known in the art
using suitable dispersing or wetting agents and
suspending agents. The sterile injectable preparations
may also be sterile injectable solutions or suspensions
in non-toxic parenterally-acceptable diluents or
15 solvents, for example, as solutions in 1,3-butanediol.
Among the acceptable vehicles and solvents that may be
employed are water, Ringer's solution and isotonic
sodium chloride solution. In addition, sterile, fixed
oils are conventionally employed as solvents or
20 suspending mediums. For this purpose, any bland fixed
oil may be employed including synthetic mono- or di-
glycerides. Fatty acids such as oleic acid and its
glyceride derivatives, including olive oil and castor
oil, especially in their polyoxyethylated versions, are
25 useful in the preparation of injectables. These oil
solutions or suspensions may also contain long-chain
alcohol diluents or dispersants.

The compounds of this invention may also be
administered rectally in the form of suppositories.
30 These compositions can be prepared by mixing the drug
with a suitable non-irritating excipient which is solid
at room temperature, but liquid at rectal temperature
and, therefore, will melt in the rectum to release the
drug. Such materials include cocoa butter, beeswax and
35 polyethylene glycols.

The compounds of this invention may also be administered topically, especially when the conditions addressed for treatment involve areas or organs readily accessible by topical application, including
5 neurological disorders of the eye, the skin, or the lower intestinal tract. Suitable topical formulations are readily prepared for each of these areas.

For topical application to the eye, or ophthalmic use, the compounds can be formulated as micronized
10 suspensions in isotonic, pH adjusted sterile saline, or, preferably, as solutions in isotonic, pH adjusted sterile saline, either with or without a preservative such as benzylalkonium chloride. Alternatively for the ophthalmic uses the compounds may be formulated in an
15 ointment such as petrolatum.

For topical application to the skin, the compounds can be formulated in a suitable ointment containing the compound suspended or dissolved in, for example, a mixture with one or more of the following: mineral oil,
20 liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene polyoxypropylene compound, emulsifying wax and water. Alternatively, the compounds can be formulated in a suitable lotion or cream containing the active compound suspended or dissolved in, for example,
25 a mixture of one or more of the following: mineral oil, sorbitan monostearate, polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

Topical application for the lower intestinal tract
30 an be effected in a rectal suppository formulation (see below) or in a suitable enema formulation.

Dosage levels on the order of about 0.1 mg to about 10,000 mg of the active ingredient compound are useful in the treatment of the above conditions, with preferred
35 levels of about 0.1 mg to about 1,000 mg. It is

understood, however, that a specific dose level for any particular patient will vary depending upon a variety of factors, including the activity of the specific compound employed; the age, body weight, general health, sex and diet of the patient; the time of administration; the rate of excretion; drug combination; the severity of the particular disease or disorder being treated; and the form of administration. Typically, *in vitro* dosage-effect results provide useful guidance on the proper doses for patient administration. Studies in animal models are also helpful. The considerations for determining the proper dose levels are well known in the art.

The specific dose may be calculated according to considerations of body weight, body surface area or organ size. Further refinement of the calculations necessary to determine the appropriate dosage for treatment involving each of the above mentioned formulations is routinely made by those of ordinary skill in the art and is within the ambit of tasks routinely performed, especially in light of the dosage information and assays disclosed herein. Appropriate dosages may be determined using established assays in conjunction with appropriate dose-response data. One skilled in the art will appreciate that the dosage used in localized formulations of the invention normally will be smaller as compared to that used in a systemic injection or oral administration.

To effectively treat alopecia or promote hair growth, the compounds used in the inventive methods and pharmaceutical compositions must readily affect the targeted areas. For these purposes, the compounds are preferably administered topically to the skin.

The compounds can be administered with other hair revitalizing agents. Specific dose levels for the other

hair revitalizing agents will depend upon the factors previously stated and the effectiveness of the drug combination. Other routes of administration known in the pharmaceutical art are also contemplated by this invention.

Specific embodiments of the inventive compounds are presented in Tables I and II. The present invention contemplates employing the compounds of Tables I and II, below, for use in compositions and methods to prevent and/or treat a neurological disorder in an animal, for use in compositions and methods to treat alopecia and promote hair growth in an animal, for use in compositions and methods to treat a vision disorder, improve vision, treat memory impairment, and enhance memory performance, and for use in compositions and methods to treat a sensorineural hearing loss in an animal, and all the other uses suggested in this specification.




TABLE I

No.	X ₁	X ₂	D	R
1	O	O	bond	Naphthyl
2	O	O	bond	2-(Phenyl)phenyl
3	O	O	bond	4-Trifluoromethylphenyl
4	S	O	methyl	Phenyl
5	O	O	hexyl	Hydrogen
6	O	O	bond	2-(Ethyl)phenyl
7	S	O	propyl	Phenyl
8	S	O	ethyl	Phenyl
9	O	O	heptyl	Hydrogen
10	O	O	octyl	Hydrogen
11	S	O	pentyl	3-Pyridyl
12	O	O	propyl	Phenyl

	No.	X ₁	X ₂	D	R
	13	O	O	bond	3-(Hydroxy)phenyl
	14	O	O	bond	4-(tert-butyl)phenyl
	15	O	O	bond	2-(Prop-2-enyl)phenyl
	16	O	O	bond	3-(Ethoxy)phenyl
5	17	S	O	bond	Cyclopentyl
	18	S	O	bond	Quinoliny
	19	O	O	hexyl	Phenyl
	20	O	O	ethyl	Phenyl
	21	O	O	bond	Cyclopentyl
10	22	S	S	bond	2-thienyl
	23	O	S	bond	2-thienyl
	24	O	O	bond	2-oxazolyl
	25	S	O	bond	2-furyl
	26	O	NH	bond	3-furyl
15	27	O	NH	hexyl	4-furyl
	28	O	S	bond	Adamantyl
	29	S	N-CN	bond	Carbazole
	30	O	N-NO ₂	bond	Isoquinoline
	31	NH	NH	methyl	3-Pyridinyl
20	32	O	NCH ₃	hexyl	Hydrogen
	33	NOH	O	bond	2-Thiazolyl
	34	NOCH ₃	S	bond	4-(tert-butyl)phenyl
	35	O	S	bond	Cyclohexyl
	36	O	O	bond	Phenyl
25	37	S	O	bond	Phenyl

Preferred compounds of TABLE I are named as follows:

(7aS)-2-(1-Naphthyl)perhydropyrrolo[1,2-c]imidazole-1,3-dione (Compound 1),

30 (7aS)-2-(2'-Phenyl)phenylperhydropyrrolo[1,2-c]imidazole-1,3-dione (Compound 2),

(7aS)-2-(4-(Trifluoromethyl)phenyl)perhydropyrrolo[1,2-c]imidazole-1,3-dione (Compound 3),

35 2-benzyl-3-thioxo-2,5,6,7,7a-pentahydro-2-azapyrrolizin-1-one (Compound 4),

2-hexyl-2,5,6,7,7a-pentahydro-2-azapyrrolizine-1,3-dione (Compound 5),

2-(2-ethyl)phenyl-2,5,6,7,7a-pentahydro-2-azapyrrolizin-1,3-dione (Compound 6),


2-(3-phenylpropyl)-3-thioxo-2,5,6,7,7a-pentahydro-2-azapyrrolizin-1-one (Compound 7),

2-(2-phenylethyl)-3-thioxo-2,5,6,7,7a-pentahydro-2-azapyrrolizin-1-one (Compound 8),

5 (7aS)-2-Cyclohexyl-3-thioxoperhydropyrrolo [1,2-c]imidazole-1-one (Compound 35),

2-Phenyl-2,5,6,7,7a-pentahydro-2-azapyrrolizine-1,3-dione (Compound 36), and

10 2-phenyl-3-thioxo-2,5,6,7,7a-pentahydro-2-azapyrrolizin-1-one (Compound 37).



(II)

TABLE II

No.	X ₁	X ₂	D	R
38	O	O	methyl	Phenyl
39	S	O	methyl	Phenyl
40	S	O	ethyl	Phenyl
41	O	O	heptyl	Hydrogen
42	O	O	octyl	Hydrogen
43	S	O	propyl	Phenyl
44	O	O	hexyl	Hydrogen
45	O	O	bond	Cyclohexyl
46	O	O	ethyl	Phenyl
47	S	O	heptyl	Hydrogen
48	O	O	octyl	Hydrogen
49	S	O	pentyl	3-Pyridyl
50	O	O	propyl	Phenyl
51	O	O	bond	3-(Phenoxy)phenyl
52	O	O	bond	4-(tert-butyl)phenyl
53	O	O	bond	2-(Prop-2-enyl)phenyl
54	O	O	bond	3-(Ethoxy)phenyl
55	S	O	bond	Cyclopentyl
56	S	O	bond	Quinolinyl

	No.	X ₁	X ₂	D	R
	57	O	O	hexyl	Phenyl
	58	O	O	ethyl	Phenyl
	59	O	O	bond	Cyclopentyl
5	60	S	S	bond	2-thienyl
	61	O	S	bond	2-thienyl
	62	O	NH	bond	2-oxazolyl
	63	S	O	bond	2-furyl
	64	O	O	bond	3-furyl
	65	S	NH	hexyl	4-furyl
10	66	O	N-CN	bond	Adamantyl
	67	S	N-NO ₂	bond	Carbazole
	68	O	S	bond	Adamantyl
	69	S	NC ₃ H ₇	bond	2-Pyrazolyl
	70	NOH	O	hexyl	Hydrogen
15	71	NOCH ₃	O	bond	Cyclopentyl
	72	O	O	bond	Phenyl
	73	S	O	bond	Phenyl
	74	O	O	butyl	Hydrogen

20 Preferred compounds of TABLE II are named as follows:

2-Benzyl-2,5,6,7,8,8a-hexahydro-2-azaindolizine-1,3-dione (Compound 38),

25 2-benzyl-3-thioxo-2,5,6,7,8,8a-hexahydro-2-azaindolizin-1-one (Compound 39),

2-(2-phenylethyl)-3-thioxo-2,5,6,7,8,8a-hexahydro-2-azaindolizin-1-one (Compound 40),

2-Heptyl-2,5,6,7,8,8a-hexahydro-2-azaindolizine-1,3-dione (Compound 41),

30 2-Octyl-2,5,6,7,8,8a-hexahydro-2-azaindolizine-1,3-dione (Compound 42),

2-(3-phenylpropyl)-3-thioxo-2,5,6,7,8,8a-hexahydro-2-azaindolizin-1-one (Compound 43),

35 2-hexyl-2,5,6,7,8,8a-hexahydro-2-azaindolizine-1,3-dione (Compound 44),

2-Cyclohexyl-2,5,6,7,8,8a-hexahydro-2-azaindolizine-1,3-dione (Compound 45),

2-phenyl-2,5,6,7,8,8a-hexahydro-2-azaindolizine-1,3-dione (Compound 72),

2-phenyl-3-thioxo-2,5,6,7,8,8a-hexahydro-2-azaindolizine-1-one (Compound 73), and

5 2-butyl-2,5,6,7,8,8a-hexahydro-2-azaindolizine-1,3-dione (Compound 74).

Other compounds which are hydantoin derivative compounds within the scope of the present invention are those compounds which may possess immunosuppressive, non-immunosuppressive, or other activities as long as they also are useful in treating a disease. In particular, other hydantoin derivative compounds falling within the scope of the present invention are useful for preventing and/or treating neurological disorders, including physically damaged nerves and neurodegenerative diseases; in treating alopecia and promoting hair growth; in treating vision disorders and/or improving vision; in treating memory impairment and/or enhancing memory performance; and/or in treating sensorineural hearing loss.

Pharmaceutical Compositions of the Present Invention

The present invention relates to a pharmaceutical composition comprising:

- 25 (i) an effective amount of a hydantoin derivative compound of formulae III or IV; and
- (ii) a pharmaceutically acceptable carrier.

The present invention also relates to a pharmaceutical composition comprising:

- 30 (i) an effective amount of a hydantoin derivative compound of formulae III or IV for treating neurodegenerative diseases, neurological disorders, and nerve damage, or promoting nerve

growth in animals; and

(ii) a pharmaceutically acceptable carrier.

The present invention also relates to a pharmaceutical composition comprising:

5 (i) an effective amount of a hydantoin derivative compound of formulae III or IV for treating alopecia or promoting hair growth in an animal; and

(ii) a pharmaceutically acceptable carrier.

10 The present invention also relates to a pharmaceutical composition comprising:

(i) an effective amount of a hydantoin derivative compound of formulae III or IV for treating a vision disorder, improving vision, treating
15 memory impairment, or enhancing memory performance in an animal; and

(ii) a pharmaceutically acceptable carrier.

The present invention also relates to a pharmaceutical composition comprising:

20 (i) an effective amount of a hydantoin derivative compound of formulae III or IV for treating sensorineural hearing loss in an animal; and

(ii) a pharmaceutically acceptable carrier.

In another preferred embodiment, the novel compounds
25 of the present invention can be administered in pharmaceutical compositions additionally containing other neurotrophic agents such as neurotrophic growth factor, brain derived growth factor, glial derived growth factor, ciliary neurotrophic factor, insulin growth factor and
30 active truncated derivatives thereof, acidic fibroblast growth factor, basic fibroblast growth factor, platelet-

derived growth factors, neurotrophin-3 and neurotrophin 4/5. The dosage level of other neurotrophic drugs will depend upon the factors previously stated and the neurotrophic effectiveness of the drug combination.

5

Methods of the Present Invention

10 The present invention relates to the use of any of the compounds seen in Tables III and IV, any of the other compounds described above, and other compounds not specifically mentioned or described herein, in the preparation of a medicament for the treatment of a disease. In a preferred embodiment, the disease to be treated includes peripheral neuropathy caused by physical injury or disease state, physical damage to the brain, 15 physical damage to the spinal cord, stroke associated with brain damage, Alzheimer's Disease, Parkinson's Disease, amyotrophic lateral sclerosis, and Huntington's Disease.

20 The present invention also relates to the use of any of the hydantoin derivative compounds described herein for the treatment of a disease. In particular, the disease to be treated includes the above-mentioned neuropathies, neurological disorders, and neurological damage.

25 The present invention also relates to a method for treating alopecia or promoting hair growth in an animal, which comprises administering to said animal an effective amount of a hydantoin derivative compound. The present invention also relates to using the inventive compounds and compositions in the preparation of a medicament for 30 the treatment of alopecia or promoting hair growth in an animal.

The inventive method is particularly useful for

treating male pattern alopecia, alopecia senilis,
alopecia areata, alopecia resulting from skin lesions or
tumors, alopecia resulting from cancer therapy such as
chemotherapy and radiation, and alopecia resulting from
5 systematic disorders such as nutritional disorders and
internal secretion disorders.

The present invention also relates to a method for
treating a vision disorder, improving vision, treating
memory impairment, or enhancing memory performance in an
10 animal, which comprises administering to said animal an
effective amount of a hydantoin derivative compound. The
present invention also relates to using the inventive
compounds and compositions in the preparation of a
medicament for the treatment of a vision disorder,
15 improving vision, treating memory impairment, or
enhancing memory performance.

The inventive methods are particularly useful for
treating various eye disorders including, but not limited
to visual disorders, diseases, injuries, and
20 complications, genetic disorders; disorders associated
with aging or degenerative vision diseases; vision
disorders correlating to physical injury to the eye,
head, or other parts of the body resulting from external
forces; disorders resulting from environmental factors;
25 disorders resulting from a broad range of diseases; and
combinations of any of the above.

In particular, the compositions and methods of the
present invention are useful for improving vision, or
correcting, treating, or preventing visual (ocular)
30 impairment or dysfunction of the visual system, including
permanent and temporary visual impairment, without
limitation. The present invention is also useful in
preventing and treating ophthalmologic diseases and
disorders, treating damaged and injured eyes, and

preventing and treating diseases, disorders, and injuries which result in vision deficiency, vision loss, or reduced capacity to see or process images, and the symptoms and complications resulting from same. The eye diseases and disorders which may be treated or prevented by the compositions and methods of the present invention are not limited with regard to the cause of said diseases or disorders. Accordingly, said compositions and methods are applicable whether the disease or disorder is caused by genetic or environmental factors, as well as any other influences. The compositions and methods of the present invention are particularly useful for eye problems or vision loss or deficiency associated with all of the following, without limitation: aging, cellular or physiological degeneration, central nervous system or neurological disorder, vascular defects, muscular defects, and exposure to adverse environmental conditions or substances.

The compositions and methods of the present invention are particularly useful in correcting, treating, or improving visual impairment, without limitation. Visual impairment in varying degrees occurs in the presence of a deviation from normal in one or more functions of the eye, including (1) visual acuity for objects at distance and near; (2) visual fields; and (3) ocular motility without diplopia. See *Physicians' Desk Reference (PDR) for Ophthalmology, 16th Edition, 6:47* (1988). Vision is imperfect without the coordinated function of all three. *Id.*

Said compositions and methods of use are also useful in correcting, treating, or improving other ocular functions including, without limitation, color perception, adaptation to light and dark, accommodation, metamorphopsia, and binocular vision. The compositions

and methods of use are particularly useful in treating, correcting, or preventing ocular disturbances including, without limitation, paresis of accommodation, iridoplegia, entropion, ectropion, epiphora,
5 lagophthalmos, scarring, vitreous opacities, non-reactive pupil, light scattering disturbances of the cornea or other media, and permanent deformities of the orbit.

The compositions and methods of use of the present invention are also highly useful in improving vision and
10 treating vision loss. Vision loss ranging from slight loss to absolute loss may be treated or prevented using said compositions and methods of use. Vision may be improved by the treatment of eye disorders, diseases, and injuries using the compositions and methods of the
15 invention. However, improvements in vision using the compositions and methods of use are not so limited, and may occur in the absence of any such disorder, disease, or injury.

The present invention also relates to a method for
20 treating a sensorineural hearing loss in an animal, which comprises administering to said animal an effective amount of a hydantoin derivative compound. The present invention also relates to using the inventive compounds and compositions in the preparation of a medicament for
25 the treatment of sensorineural hearing loss.

It is further contemplated that administration of an inventive compound will protect hair cells and spiral ganglion neurons from traumatic damage, for example damage caused by noise trauma, acute or chronic treatment
30 with cisplatin and aminoglycoside antibiotics or from damage resulting from a lack of neurotrophic factors resulting from interruption of transport of the factors from the axon to the cell body. Such treatment is expected to allow hair cells and/or auditory neurons to

tolerate intermittent insults from either environmental noise trauma or treatment with ototoxins, and to slow down, prevent or reverse the progressive degeneration of the auditory neurons and hair cells which is responsible for hearing loss in pathological conditions such as presbycusis (age-related hearing loss), inherited sensorineural degeneration, and post-idiopathic hearing losses and to preserve the functional integrity of the inner ear. Such treatment will also support the auditory neurons for better and longer performance of cochlear implants.

It is understood, however, that a specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, rate of excretion, drug combination, and the severity of the particular disease or disorder being treated and form of administration.

20 K. Test Procedure

The binding to FBKP12 and inhibition of the peptidyl-prolyl isomerase (rotamase) activity of the compounds used in the inventive methods and pharmaceutical compositions can be evaluated by known methods described in the literature (Harding et al., *Nature*, 1989, 341:758-760; Holt et al. *J. Am. Chem. Soc.*, 115:9923-9938). These values are obtained as apparent K_i 's and are presented for representative compounds in TABLE III.

30 The *cis-trans* isomerization of an alanine-proline bond in a model substrate, N-succinyl-Ala-Ala-Pro-Phe-p-nitroanilide, is monitored spectrophotometrically in a chymotrypsin-coupled assay, which releases *para*-

nitroanilide from the *trans* form of the substrate. The inhibition of this reaction caused by the addition of different concentrations of inhibitor is determined, and the data is analyzed as a change in first-order rate constant as a function of inhibitor concentration to yield the apparent K_i values.

In a plastic cuvette are added 950 μ L of ice cold assay buffer (25 mM HEPES, pH 7.8, 100 mM NaCl), 10 μ L of FKBP (2.5 mM in 10 mM Tris-Cl pH 7.5, 100 mM NaCl, 1 mM dithiothreitol), 25 μ L of chymotrypsin (50 mg/mL in 1 mM HCl) and 10 μ L of test compound at various concentrations in dimethyl sulfoxide. The reaction is initiated by the addition of 5 μ L of substrate (succinyl-Ala-Phe-Pro-Phe-*para*-nitroanilide, 5 mg/mL in 2.35 mM LiCl in trifluoroethanol).

The absorbance at 390 nm versus time is monitored for 90 seconds using a spectrophotometer and the rate constants are determined from the absorbance versus time data files.

TABLE III

In Vitro Test Results

Compound	K_i (nM)
1	>10,000
6	>5,000
35	>10,000

MPTP Model of Parkinson's Disease in Mice

MPTP lesioning of dopaminergic neurons in mice was used as an animal model of Parkinson's Disease. Four week old male CD1 white mice were dosed i.p. with 30 mg/kg of MPTP for 5 days. Inventive compounds (4 mg/kg), or vehicle, were administered s.c. along with the MPTP for 5 days, as well as for an additional 5 days following

cessation of MPTP treatment (Table IV). Inventive compounds (10 mg/kg), or vehicle, were also administered p.o. along with the MPTP for 5 days, as well as for an additional 5 days following cessation of MPTP treatment (Table V). At 18 days following MPTP treatment, the animals were sacrificed and the striata were dissected and homogenized. Immunostaining was performed on sagittal and coronal brain sections using anti-tyrosine hydroxylase Ig to quantitate survival and recovery of dopaminergic neurons. In animals treated with MPTP and vehicle, a substantial loss of functional dopaminergic terminals was observed as compared to non-lesioned animals. In another protocol, test compounds were administered only subsequent to MPTP-induced lesioning. Thus, after animals were treated with MPTP for 5 days, an additional 3 days passed before beginning oral drug treatment on day 8. Animals were treated with inventive compounds (0.4 mg/kg), administered orally, once a day for 5 days total. On day 18, the animals were sacrificed and analyzed as described above. Tables IV and V present the percent recovery of dopaminergic neurons in the first (concurrent dosing) paradigms in animals receiving hydantoin derivative compounds.

Tables IV and V, below, show the remarkable neuroregenerative effects of the inventive hydantoin derivative compounds illustrating the neurotrophic capability of these compounds as a class showing that lesioned animals receiving the hydantoin derivative compounds provide a remarkable recovery of TH-stained dopaminergic neurons.

TABLE IV

% Recovery

Compound 1	32.20 %
------------	---------

TABLE V

		<u>% Recovery</u>
	Compound 2	38.40 %
	Compound 3	78.80 %
5	Compound 4	49.30 %
	Compound 7	37.50 %
	Compound 8	42.50 %
	Compound 35	24.40 %
	Compound 38	31.40 %
10	Compound 39	11.20 %
	Compound 40	11.00 %
	Compound 41	8.80 %
	Compound 42	17.40 %
	Compound 43	40.10 %
15	Compound 72	37.20 %
	Compound 74	46.40 %

Percent striatal innervation density was quantitated in brain sections with an anti-tyrosine hydroxylase immunoglobulin, which is indicative of functional dopaminergic neurons. The striatal innervation density of 23% for animals pretreated with only a vehicle and administered a vehicle orally during treatment, is indicative of normal non-lesioned striatal tissue. Striatal innervation density is reduced to 5% for animals pretreated with MPTP and administered a vehicle orally during treatment, and is indicative of MPTP-induced lesioning. Surprisingly, striatal innervation density is increased 8%-13% for animals pretreated with MPTP and administered 0.4 mg/kg orally during treatment, indicating

substantial neuronal regeneration after induction of MPTP-derived lesions.

Synthesis of Compounds of the Invention


5 The compounds for use in the methods and compositions of the invention may be readily prepared by standard techniques of organic chemistry, utilizing the general synthetic pathways depicted below.

10 In the preparation of the compounds of the invention, one skilled in the art will understand that one may need to protect or block various reactive functionalities on the starting compounds or intermediates while a desired reaction is carried out on other portions of the molecule. After the desired reactions are complete, or at any desired time, normally
15 such protecting groups will be removed by, for example, hydrolytic or hydrogenolytic means. Such protection and deprotection steps are conventional in organic chemistry. One skilled in the art is referred to "Protective Groups in Organic Chemistry," McOmie, ed., Plenum Press, New
20 York, New York; and "Protective Groups in Organic Synthesis," Greene, ed., John Wiley & Sons, New York, N.Y. (1981) for the teaching of protective groups which may be useful in the preparation of compounds of the present invention.

25 The product and intermediates may be isolated or purified using one or more standard purification techniques, including, for example, one or more of simple solvent evaporation, recrystallization, distillation, sublimation, filtration, chromatography, including thin-
30 layer chromatography, HPLC (e.g. reverse phase HPLC), column chromatography, flash chromatography, radial chromatography, trituration, and the like.

 The compounds of the present invention may be

prepared by reacting amino acids with isocyanates and isothiocyanates, as shown in the general method of Scheme I:



5

Scheme I

EXAMPLES

The following examples are illustrative of the present invention and are not intended to be limitations thereon. Unless otherwise indicated, all percentages are based upon 100% by weight of the final composition.

EXAMPLE 1

Synthesis of 2-phenyl-2,5,6,7,8,8a-hexahydro-2-azaindolizine-1,3-dione

A solution of L-pipecolic acid methyl ester (1.0 g; 3.58 mmol), phenylisocyanate (0.42 ml; 3.93 mmol), and triethylamine (1.64 ml; 11.8 mmol) in methylene chloride (15 ml) was stirred overnight at room temperature. The solvent was removed under vacuum, the residue was dissolved in 50 ml of 50% ethyl acetate in hexane, and the organic phase was washed with water and brine, dried, and concentrated. The residue was purified on a flash column, eluting with 40% ethyl acetate in hexane, to obtain the product as a white solid, ¹H NMR (MeOH, 400 MHz): δ 1.47-1.57 (cm, 3H); 1.62 (bd, 1H, J=9.99); 2.01 (bd, 1H, J=10.36); 2.20 (bd, 1H, J=9.88); 2.96 (t, 1H,

J=9.35); 4.06-4.17 (m, 2H); 7.35-7.40 (m, 3H); 7.45-7.48 (m, 2H). Anal. Calcd. for: C, 67.81; H, 6.13; N, 12.17. Found: C, 67.93; H, 6.15; N, 12.21. Mp = 186.1 - 186.7 °C.

5

EXAMPLE 2

Synthesis of 2-(2-phenylethyl)-3-thioxo-2,5,6,7,7a-pentahydro-2-azapyrrolizin-1-one

A solution of L-proline methyl ester (1.0 g; 6.03 mmol), phenethylisothiocyanate (1.08 ml; 6.63 mmol), and triethylamine (2 ml; 15.07 mmol) in 15 ml of methylene chloride was stirred overnight at room temperature. The reaction was worked up as described previously, and the product purified to deliver a white solid, ¹H NMR (CDCl₃, 400 MHz): δ 1.56-1.62 (m, 1H); 2.10-2.26 (m, 3H); 2.96-3.01 (m, 2H); 3.52-3.58 (m, 1H); 3.92-3.99 (m, 3H); 4.00-4.12 (m, 1H); 7.19-7.31 (m, 5H). Anal. Calcd. for: C, 64.59; H, 6.19; N, 10.76; S, 12.32. Found: C, 64.69; H, 6.13; N, 10.71; S, 12.44. Mp = 61.2-63.5 °C

EXAMPLE 3

2-(3-phenylpropyl)-3-thioxo-2,5,6,7,7a-pentahydro-2-azapyrrolizin-1-one

This example was prepared according to the method previously described. ¹H NMR (CDCl₃, 400 MHz): δ 1.60-1.63 (m, 1H); 2.00-2.06 (m, 2H); 2.12-2.18 (m, 1H); 2.20-2.26 (m, 2H); 2.66 (t, 2H, J=7.65); 3.51-3.57 (m, 1H); 3.80-3.85 (m, 2H); 3.90-3.97 (m, 1H); 4.07-4.12 (m, 1H); 7.15-7.25 (m, 5H). Anal. Calcd. for: C, 65.66; H, 6.61; N, 10.21; S, 11.69. Found: C, 65.66; H, 6.56; N, 10.14; S, 11.71.

30

EXAMPLE 4

(7aS)-2-Cyclohexyl-3-thioxoperhydropyrrolo[1,2-climidazole-1-one

This example was prepared according to the method
5 previously described. ^1H NMR (CDCl_3 , 400 MHz): δ 1.12-
1.41 (m, 3H); 1.59-1.65 (m, 4H); 1.69-1.81 (m, 2H); 2.10-
2.21 (m, 2H); 2.25-2.38 (m, 3H); 3.61-3.66 (m, 1H); 3.90-
3.98 (m, 1H); 4.02-4.13 (m, 1H); 4.44-4.51 (m, 1H). Anal.
Calcd. for $\text{C}_{12}\text{H}_{18}\text{N}_2\text{OS}$: C, 60.47; H, 7.61; N, 11.75; S,
10 13.45. Found: C, 60.52; H, 7.60; N, 11.73; S, 13.27.

EXAMPLE 5

(7aS)-2-(2'-Phenyl)phenylperhydropyrrolo[1,2-climidazole-
1,3-dione

This example was prepared according to the method
15 previously described. ^1H NMR (CDCl_3 , 400 MHz): δ 1.00-1.16
(m, 1H); 1.58-2.25 (cm, 3H); 3.12-3.22 (m, 1H); 3.57-3.72
(m, 1H); 3.92-4.04 (m, 1H); 7.20-7.52 (m, 9H). Anal.
Calcd. for $\text{C}_{18}\text{H}_{16}\text{N}_2\text{O}_2$ -0.05 EtOAc: C, 73.67; H, 5.57; N,
9.44. Found: C, 73.50; H, 5.87; N, 9.74.

EXAMPLE 6

(7aS)-2-(4-(Trifluoromethyl)phenyl)perhydropyrrolo[1,2-
climidazole-1,3-dione

This example was prepared according to the method
previously described. ^1H NMR (CDCl_3 , 400 MHz): δ 1.82-
25 1.99 (m, 1H); 2.14-2.22 (m, 2H); 2.34-2.39 (m, 1H); 3.34-
3.40 (m, 1H); 3.77-3.84 (m, 1H); 4.24-4.29 (m, 1H); 7.67
(d, 2H, $J = 8.35$); 7.72 (d, 2H, $J = 8.60$). Anal. Calcd.
for $\text{C}_{13}\text{H}_{11}\text{N}_2\text{O}_2\text{F}$: C, 54.93; H, 3.90; N, 9.86. Found: C,
55.45; H, 4.26; N, 9.87.

EXAMPLE 7

(7aS)-2-(1-Naphthyl)perhydropyrrolo[1,2-climidazole-1,3-dione

This example was prepared according to the method
5 previously described. ¹H NMR (CDCl₃, 300 MHz): δ 1.94-
2.03 (m, 1H); 2.05-2.33 (m, 2H); 2.39-2.51 (m, 1H); 3.38-
3.44 (m, 1H); 3.83-4.09 (m, 1H); 4.41 (dt, 1H, J = 7.45);
7.38-7.96 (m, 2H). Anal. Calcd. for C₁₆H₁₄N₂O₂: C, 72.17;
H, 5.30; N, 10.52. Found: C, 72.05; H, 5.32; N, 10.51.

10

EXAMPLE 8

2-Phenyl-2,5,6,7,7a-pentahydro-2-azapyrrolizine-1,3-dione

This example was prepared according to the method
previously described. ¹H NMR (CDCl₃, 400 MHz): δ 1.78 (m,
1H); 2.10 (m, 2H); 2.26 (m, 1H); 3.28 (m, 1H); 3.73 (q,
15 1H); 4.17 (t, 2H); 7.31 (m, 3H); 7.38 (m, 2H).

EXAMPLE 9

2-benzyl-2,5,6,7,8,8a-hexahydro-2-azaindolizine-1,3-dione

This example was prepared according to the method
previously described. ¹H NMR (MeOH, 400 MHz): δ 1.26-1.39
20 (m, 2H); 1.50-1.57 (m, 1H); 1.7 (bd, 1H, J=10.03); 1.95
(bd, 1H, J=13.56); 2.13 (bd, 1H, J=9.57); 2.89 (td, 1H,
J=9.44, 3.52); 3.93 (dd, 1H, j=12.07, 4.26); 4.04-4.09
(m, 1H); 4.61 (s, 2H); 7.24-7.31 (m, 5H). Anal. Calcd.
for: C, 68.83; H, 6.60; N, 11.47. Found: C, 68.90; H,
25 6.61; N, 11.48.

EXAMPLE 10

2-benzyl-3-thioxo-2,5,6,7,7a-pentahydro-2-azapyrrolizin-
1-one

This example was prepared according to the method
30 previously described. ¹H NMR (MeOH, 400 MHz): δ 1.60-1.66

(m, 1H); 2.20-2.26 (m, 3H); 3.50-3.56 (m, 1H); 3.86-3.93 (m, 1H); 4.33-4.37 (m, 1H); 4.95 (d, 2H, J=14.81); 7.20-7.29 (m, 3H); 7.30-7.37 (m, 2H). Anal. Calcd. for: C, 63.39; H, 5.73; N, 11.37; S, 13.02. Found: C, 63.60; H, 5.90; N, 11.21; S, 12.92.

EXAMPLE 11

2-benzyl-3-thioxo-2,5,6,7,8,8a-hexahydro-2-azaindolizine-1-one

This example was prepared according to the method previously described. ¹H NMR (MeOH, 400 MHz): δ 1.30-1.33 (m, 1H); 1.42-1.45 (m, 1H); 1.69 (q, 1H, J=9.55); 1.82 (bd, 1H, J=13.38); 1.95 (bd, 1H, J=13.55); 2.16 (bd, 1H, J=12.91); 3.09 (td, 1H, J=3.27, 13.11); 4.07 (dd, 1H, J=7.18, 4.13); 4.76 (dd, 1H, J=9.46, 3.46); 4.97 (d, 2H); 7.22-7.29 (m, 3H); 7.36-7.38 (m, 2H). Anal. Calcd. for: C, 64.59; H, 6.19; N, 10.76; S, 12.32. Found: C, 64.69; H, 6.22; N, 10.83; S, 12.32.

EXAMPLE 12

2-(2-phenylethyl)-3-thioxo-2,5,6,7,8,8a-hexahydro-2-azaindolizine-1-one

This example was prepared according to the method previously described. ¹H NMR (CDCl₃, 400 MHz): δ 1.22-1.26 (m, 1H); 1.44-1.49 (m, 2H); 1.82 (bd, 1H, J=10.00); 1.95 (bd, 1H, J=11.04); 2.18 (bd, 1H, J=10.11); 2.95-3.03 (m, 3H); 3.76 (dd, 1H, J=4.34, 12.14); 4.01-4.06 (m, 2H); 4.83 (bd, 1H, J=13.32); 7.19-7.29 (m, 5H). Anal. Calcd. - with 0.20 H₂O for: C, 64.81; H, 6.67; N, 10.08; S, 11.53. Found: C, 64.64; H, 6.60; N, 9.92; S, 11.57.

EXAMPLE 13

2-heptyl-2,5,6,7,8,8a-hexahydro-2-azaindolizine-1,3-dione

This example was prepared according to the method

previously described. ^1H NMR (MeOH, 400 MHz): δ 0.87-0.91 (m, 4H); 1.30 (bs, 9H); 1.53-1.59 (m, 3H); 1.73 (bd, 1H, $J=13.17$); 1.94 (bd, 1H, $J=10.43$); 2.13 (bd, 1H, $J=12.69$); 2.88 (td, 1H, $J=3.50, 9.57$); 3.30 (s, 1H); 3.43-3.47 (m, 1H); 3.89 (dd, 1H, $J=4.20, 11.94$); 4.06 (dd, 1H, $J=3.85, 12.85$). Anal. Calcd. for: C, 66.63; H, 9.59; N, 11.10. Found: C, 66.37; H, 9.56; N, 10.99.

EXAMPLE 14

2-octyl-2,5,6,7,8,8a-hexahydro-2-azaindolizine-1,3-dione

This example was prepared according to the method previously described. ^1H NMR (CDCl_3 , 400 MHz): δ 0.87 (bs, 3H); 1.28 (bs, 11H); 1.36-1.50 (m, 2H); 1.59 (bs, 2H); 1.97 (bd, 1H, $J=11.28$); 2.18 (bd, 1H, $J=13.16$); 2.22 (bd, 1H, $J=13.12$); 2.81 (td, 1H, $J=3.45, 12.86$); 3.47 (t, 2H, $J=7.39$); 3.72 (dd, 1H, $J=4.24, 11.96$); 4.17 (dd, 1H, $J=4.71, 13.31$). Anal. Calcd. for: C, 67.63; H, 9.84; N, 10.52. Found: C, 67.89; H, 10.01; N, 10.42.

EXAMPLE 15

2-phenyl-3-thioxo-2,5,6,7,8,8a-hexahydro-2-azaindolizine-1-one

This example was prepared according to the method previously described. ^1H NMR (MeOH, 400 MHz): δ 1.54-1.66 (m, 3H); 1.88 (bd, 1H, $J=8.49$); 1.90-2.08 (m, 1H); 2.35 (bd, 1H, $J=11.96$); 3.10 (td, 1H, $J=3.68, 12.85$); 4.04 (dd, 1H, $J=4.48, 11.67$); 4.93 (dd, 1H, $J=1.95, 11.03$); 7.30-7.32 (m, 2H); 7.41-7.51 (m, 3H). Anal. Calcd. for: C, 63.39; H, 5.73; N, 11.37; S, 13.02. Found: C, 63.53; H, 5.82; N, 11.32; S, 13.07.

EXAMPLE 16

2-(3-phenylpropyl)-3-thioxo-2,5,6,7,8,8a-hexahydro-2-azaindolizine-1-one

This example was prepared according to the method previously described. ¹H NMR (CDCl₃, 400 MHz): δ 1.23-1.29 (m, 2H); 1.47-1.54 (m, 2H); 1.79 (bd, 1H, J=10.18); 1.82-2.08 (m, 2H); 2.22 (bd, 1H, J=13.02); 2.67 (t, 2H, J=7.61); 2.95 (td, 1H, J=3.33, 9.59); 3.71 (dd, 1H, J=4.31, 12.14); 3.85-3.89 (m, 2H); 4.82 (dd, 1H, J=4.45, 13.18); 7.14-7.28 (m, 5H). Anal. Calcd. for: C, 66.63; H, 6.99; N, 9.71; S, 11.12. Found: C, 66.75; H, 7.08; N, 9.67; S, 11.21.

EXAMPLE 17

2-butyl-2,5,6,7,8,8a-hexahydro-2-azaindolizine-1,3-dione

This example was prepared according to the method previously described. ¹H NMR (CDCl₃, 400 MHz): δ 0.92 (t, 3H, J=7.33); 1.24-1.63 (cm, 7H); 1.73 (bd, 1H, J=8.55); 1.98 (bd, 1H, J=13.44); 2.21 (bd, 1H, J=9.90); 2.81 (td, 1H, J=3.56, 12.78); 3.49 (t, 2H, J=7.34); 3.72 (dd, 1H, J=4.26, 11.97); 4.15 (dd, 1H, J=4.74, 13.33). Anal. Calcd. for: C, 62.83; H, 8.63; N, 13.32. Found: C, 62.57; H, 8.54; N, 13.17.

EXAMPLE 18

2-hexyl-2,5,6,7,8,8a-hexahydro-2-azaindolizine-1,3-dione

This example was prepared according to the method previously described. ¹H NMR (MeOH, 400 MHz): δ 0.88-0.91 (m, 3H); 1.25-1.34 (bs, 8H); 1.50-1.58 (m, 3H); 1.71-1.75 (m, 1H); 1.92-1.96 (m, 1H); 2.11-2.15 (m, 1H); 2.85-2.93 (m, 1H); 3.43-3.47 (m, 2H); 3.87-3.91 (m, 1H); 4.04-4.08 (m, 1H). Anal. Calcd. with 0.1 EtOAc for: C, 65.12; H, 9.30; N, 11.34. Found: C, 64.87; H, 9.26; N,

11.54.

EXAMPLE 19

2-hexyl-2,5,6,7,7a-pentahydro-2-azapyrrolizine-1,3-dione

This example was prepared according to the method
5 previously described. ¹H NMR (CDCl₃, 400 MHz): δ 0.85-
0.89 (bs, 3H); 1.28 (bs, 7H); 1.59 (m, 1H); 1.63-1.73 (m,
1H); 2.22-2.24 (m, 2H); 2.25-2.27 (m, 1H); 3.21-3.27 (m,
1H); 3.43-3.47 (m, 2H); 3.65-3.72 (m, 1H); 4.06 (dd, 1H,
J=1.20, 7.22). Anal. Calcd. for: C, 64.26; H, 8.99; N,
10 12.49. Found: C, 64.02; H, 9.00; N, 12.42.

EXAMPLE 20

2-phenyl-3-thioxo-2,5,6,7,7a-pentahydro-2-azapyrrolizin-
1-one

This example was prepared according to the method
15 previously described. ¹H NMR (CDCl₃, 400 MHz): δ 1.85-
1.93 (m, 1H); 2.21-2.27 (m, 1H); 2.31-2.43 (m, 2H); 3.63-
3.69 (m, 1H); 4.04-4.11 (m, 1H); 4.38 (dd, 1H, J=6.96,
10.39); 7.25-7.32 (m, 2H); 7.40-7.51 (m, 3H). Mp =
187.3-189.7 °C. Anal. Calcd. for: C, 62.04; H, 5.21; N,
20 12.06; S, 13.80. Found: C, 61.85; H, 5.26; N, 12.01; S,
13.97.

EXAMPLE 21

A lotion comprising the following composition may be
prepared.

25		(%)
	95% Ethanol	80.0
	a hydantoin derivative	10.0
	α-Tocopherol acetate	0.01
30	Ethylene oxide (40 mole) adducts of hardened castor oil	0.5

purified water	9.0
perfume and dye	q.s.

5 Into 95% ethanol are added a hydantoin derivative, α -tocopherol acetate, ethylene oxide (40 mole) adducts of hardened castor oil, perfume, and a dye. The resulting mixture is stirred and dissolved, and purified water is added to the mixture to obtain a transparent liquid lotion.

10 5 ml of the lotion may be applied once or twice per day to a site having marked baldness or alopecia.

EXAMPLE 22

A lotion comprising the following composition shown may be prepared.

		(%)
15	95% Ethanol	80.0
	a hydantoin derivative	0.005
	Hinokitol	0.01
	Ethylene oxide (40 mole) adducts of hardened castor oil	0.5
20	Purified water	19.0
	Perfume and dye	q.s.

25 Into 95% ethanol are added a hydantoin derivative, hinokitol, ethylene oxide (40 mole) adducts of hardened castor oil, perfume, and a dye. The resulting mixture is stirred, and purified water is added to the mixture to obtain a transparent liquid lotion.

The lotion may be applied by spraying once to 4 times per day to a site having marked baldness or alopecia.

30

EXAMPLE 23

An emulsion may be prepared from A phase and B phase

having the following compositions.

	(A phase)	(%)
	Whale wax	0.5
	Cetanol	2.0
5	Petrolatum	5.0
	Squalane	10.0
	Polyoxyethylene (10 mole) monostearate	2.0
	Sorbitan monooleate	1.0
	a hydantoin derivative	0.01
10	(B phase)	(%)
	Glycerine	10.0
	Purified water	69.0
	Perfume, dye, and preservative	q.s.

15 The A phase and the B phase are respectively heated and melted and maintained at 80°C. Both phases are then mixed and cooled under stirring to normal temperature to obtain an emulsion.

20 The emulsion may be applied by spraying once to four times per day to a site having marked baldness or alopecia.

EXAMPLE 24

A cream may be prepared from A phase and B phase having the following compositions.

	(A Phase)	(%)
25	Fluid paraffin	5.0
	Cetostearyl alcohol	5.5
	Petrolatum	5.5
	Glycerine monostearate	33.0
30	Polyoxyethylene (20 mole) 2-octyldodecyl ether	3.0

	Propylparaben	0.3
	(B Phase)	(%)
	a hydantoin derivative	0.8
	Glycerine	7.0
5	Dipropylene glycol	20.0
	Polyethylene glycol 4000	5.0
	Sodium Hexametaphosphate	0.005
	Purified water	44.895

10 The A phase is heated and melted, and maintained at 70°C. The B phase is added into the A phase and the mixture is stirred to obtain an emulsion. The emulsion is then cooled to obtain a cream.

15 The cream may be applied once to 4 times per day to a site having marked baldness or alopecia.

EXAMPLE 25

A topical liquid comprising the following composition may be prepared.

		(%)
20	Polyoxyethylene butyl ether	20.0
	Ethanol	50.0
	a hydantoin derivative	0.001
	Propylene glycol	5.0
25	Polyoxyethylene hardened castor oil derivative (ethylene oxide 80 mole adducts)	0.4
	Perfume	q.s.
	Purified water	q.s.

30 Into ethanol are added polyoxypropylene butyl ether, propylene glycol, polyoxyethylene hardened castor oil, a hydantoin derivative, and perfume. The resulting mixture

is stirred, and purified water is added to the mixture to obtain a liquid.

The liquid may be applied once to 4 times per day to a site having marked baldness or alopecia.

5

EXAMPLE 26

A shampoo comprising the following composition may be prepared.

10

	(%)
Sodium laurylsulfate	5.0
Triethanolamine laurylsulfate	5.0
Betaine lauryldimethylaminoacetate	6.0
Ethylene glycol distearate	2.0
Polyethylene glycol	5.0
a hydantoin derivative	5.0
15 Ethanol	2.0
Perfume	0.3
Purified water	69.7

15

20 Into 69.7 of purified water are added 5.0 g of sodium laurylsulfate, 5.0 g of triethanolamine laurylsulfate, 6.0 g of betaine lauryldimethyl-aminoacetate. Then a mixture obtained by adding 5.0 g of a hydantoin derivative, 5.0 g of polyethylene glycol, and 2.0 g of ethylene glycol distearate to 2.0 g of ethanol, followed by stirring, and 25 0.3 g of perfume are successively added. The resulting mixture is heated and subsequently cooled to obtain a shampoo.

The shampoo may be used on the scalp once or twice per day.

30

EXAMPLE 27

A patient is suffering from alopecia senilis. A

hydantoin derivative compound, or a pharmaceutical composition comprising the same, may be administered to the patient. Increased hair growth is expected to occur following treatment.

5

EXAMPLE 28

A patient is suffering from male pattern alopecia. A hydantoin derivative compound, or a pharmaceutical composition comprising the same, or a pharmaceutical composition comprising the same may be administered to the patient. Increased hair growth is expected to occur following treatment.

10

EXAMPLE 29

A patient is suffering from alopecia areata. A hydantoin derivative compound, or a pharmaceutical composition comprising the same, may be administered to the patient. Increased hair growth is expected to occur following treatment.

15

EXAMPLE 30

A patient is suffering from hair loss caused by skin lesions. A hydantoin derivative compound, or a pharmaceutical composition comprising the same, may be administered to the patient. Increased hair growth is expected to occur following treatment.

20

EXAMPLE 31

A patient is suffering from hair loss caused by tumors. A hydantoin derivative compound, or a pharmaceutical composition comprising the same, may be administered to the patient. Increased hair growth is expected to occur following treatment.

25

EXAMPLE 32

30

A patient is suffering from hair loss caused by a

systematic disorder, such as a nutritional disorder or an internal secretion disorder. A hydantoin derivative compound, or a pharmaceutical composition comprising the same, may be administered to the patient. Increased hair growth is expected to occur following treatment.

EXAMPLE 33

A patient is suffering from hair loss caused by chemotherapy. A hydantoin derivative compound, or a pharmaceutical composition comprising the same, may be administered to the patient. Increased hair growth is expected to occur following treatment.

EXAMPLE 34

A patient is suffering from hair loss caused by radiation. A hydantoin derivative compound, or a pharmaceutical composition comprising the same may, be administered to the patient. Increased hair growth is expected to occur following treatment.

EXAMPLE 35

A patient is suffering from a neurodegenerative disease. A hydantoin derivative compound or a pharmaceutical composition comprising the same is administered. It would be expected that the patient would improve their condition or recover.

EXAMPLE 36

A patient is suffering from a neurological disorder. A hydantoin derivative compound or pharmaceutical compositions comprising same is administered. It would be expected that the patient would improve their condition or recover.

EXAMPLE 37

A patient is suffering from stroke. A hydantoin

derivative compound or pharmaceutical compositions comprising same is administered. It would be expected that the patient would improve their condition or recover.

EXAMPLE 38

5 A patient is suffering from Parkinson's Disease. A hydantoin derivative compound or pharmaceutical compositions comprising same is administered. It would be expected that the patient would improve their condition or recover.

10 EXAMPLE 39

 A patient is suffering from Alzheimer's Disease. A hydantoin derivative compound or pharmaceutical compositions comprising same is administered. It would be expected that the patient would improve their condition or
15 recover.

EXAMPLE 40

 A patient is suffering from Huntington's Disease. A hydantoin derivative compound or pharmaceutical compositions comprising same is administered. It would be expected that the patient would improve their condition or
20 recover.

EXAMPLE 41

 A patient is suffering from a peripheral neuropathy. A hydantoin derivative compound or pharmaceutical
25 compositions comprising same is administered. It would be expected that the patient would improve their condition or recover.

EXAMPLE 42

 A patient is suffering from amyotrophic lateral
30 sclerosis. A hydantoin derivative compound or pharmaceutical compositions comprising same is

administered. It would be expected that the patient would improve their condition or recover.

EXAMPLE 43

5 A patient is suffering from a spinal injury. A hydantoin derivative compound or pharmaceutical compositions comprising same is administered. It would be expected that the patient would improve their condition or recover.

EXAMPLE 44

10 A patient is at risk of suffering from a neurodegenerative disease or neurological disorder. A hydantoin derivative compound or a pharmaceutical composition comprising the same is prophelactically administered. It would be expected that the patient would
15 be prevented from some or all of the effects of the disease or disorder, or would significantly improve their condition or recover over patients who were not pre-treated.

EXAMPLE 45

20 A patient is suffering from macular degeneration. A hydantoin derivative compound as identified above, alone or in combination with one or more other neopsic factors, or a pharmaceutical composition comprising the same, may be administered to the patient. A reduction in vision loss, prevention of vision degeneration, and/or promotion of
25 vision regeneration are/is expected to occur following treatment.

EXAMPLE 46

30 A patient is suffering from glaucoma, resulting in cupping of the optic nerve disc and damage to nerve fibers. A hydantoin derivative compound as identified above, alone or in combination with one or more other neopsic factors, or a pharmaceutical composition comprising the same, may be

administered to the patient. A reduction in vision loss, prevention of vision degeneration, and/or promotion of vision regeneration are/is expected to occur following treatment.

5

EXAMPLE 47

A patient is suffering from cataracts requiring surgery. Following surgery, a hydantoin derivative compound as identified above, alone or in combination with one or more other neoplastic factors, or a pharmaceutical composition comprising the same, may be administered to the patient. A reduction in vision loss, prevention of vision degeneration, and/or promotion of vision regeneration are/is expected to occur following treatment.

10

EXAMPLE 48

A patient is suffering from an impairment or blockage of retinal blood supply relating to diabetic retinopathy, ischemic optic neuropathy, or retinal artery or vein blockage. A hydantoin derivative compound as identified above, alone or in combination with one or more other neoplastic factors, or a pharmaceutical composition comprising the same, may be administered to the patient. A reduction in vision loss, prevention of vision degeneration, and/or promotion of vision regeneration are/is expected to occur following treatment.

15

20

25

EXAMPLE 49

A patient is suffering from a detached retina. A hydantoin derivative compound as identified above, alone or in combination with one or more other neoplastic factors, or a pharmaceutical composition comprising the same, may be administered to the patient. A reduction in vision loss, prevention of vision degeneration, and/or promotion of vision regeneration are/is expected to occur following treatment.

30

EXAMPLE 50

A patient is suffering from tissue damage caused by inflammation associated with uveitis or conjunctivitis. A hydantoin derivative compound as identified above, alone or
5 in combination with one or more other neoplastic factors, or a pharmaceutical composition comprising the same, may be administered to the patient. A reduction in vision loss, prevention of vision degeneration, and/or promotion of vision regeneration are/is expected to occur following
10 treatment.

EXAMPLE 51

A patient is suffering from photoreceptor damage caused by chronic or acute exposure to ultraviolet light. A hydantoin derivative compound as identified above, alone
15 or in combination with one or more other neoplastic factors, or a pharmaceutical composition comprising the same, may be administered to the patient. A reduction in vision loss, prevention of vision degeneration, and/or promotion of vision regeneration are/is expected to occur following
20 treatment.

EXAMPLE 52

A patient is suffering from optic neuritis. A hydantoin derivative compound as identified above, alone or
25 in combination with one or more other neoplastic factors, or a pharmaceutical composition comprising the same, may be administered to the patient. A reduction in vision loss, prevention of vision degeneration, and/or promotion of vision regeneration are/is expected to occur following
treatment.

30 EXAMPLE 53

A patient is suffering from tissue damage associated with a "dry eye" disorder. A hydantoin derivative compound

as identified above, alone or in combination with one or more other neopsic factors, or a pharmaceutical composition comprising the same, may be administered to the patient. A reduction in vision loss, prevention of vision
5 degeneration, and/or promotion of vision regeneration are/is expected to occur following treatment.

EXAMPLE 54

A patient is suffering from sensorineural hearing loss. A hydantoin derivative compound as identified
10 above, alone or in combination with one or more other factors, or a pharmaceutical composition comprising the same, may be administered to the patient. A reduction in hearing loss is expected to occur following treatment.

The invention being thus described, it will be
15 obvious that the same may be varied in many ways. Such variations are not to be regarded as a departure from the spirit and scope of the invention and all such modifications are intended to be included within the scope of the following claims.